



UNIVERSIDADE FEDERAL RURAL DE PERNAMBUCO
PRÓ-REITORIA DE PESQUISA E PÓS-GRADUAÇÃO
PROGRAMA DE PÓS-GRADUAÇÃO EM RECURSOS PESQUEIROS E
AQUICULTURA

EFEITO DA FREQUÊNCIA DO AJUSTE DO EQUILÍBRIO IÔNICO NO
DESEMPENHO ZOOTÉCNICO DO *Litopenaeus vannamei* (BOONE, 1931) EM
SISTEMA SIMBIÓTICO EM ÁGUAS OLIGOHALINAS

Valdemir Queiroz de Oliveira

Dissertação apresentada ao Programa de Pós-Graduação em Recursos Pesqueiros e Aquicultura da Universidade Federal Rural de Pernambuco como exigência para obtenção do título de mestre.

Prof. Dr. Alfredo Olivera Gálvez

Orientador

Prof. Dr. Luis Otavio Brito da Silva

Coorientador

Recife

2021

Dados Internacionais de Catalogação na Publicação
Universidade Federal Rural de Pernambuco
Sistema Integrado de Bibliotecas
Gerada automaticamente, mediante os dados fornecidos pelo(a) autor(a)

O48e

Oliveira, Valdemir Queiroz de

Efeito da frequência do ajuste do equilíbrio iônico no desempenho zootécnico do *Litopenaeus vannamei* (BOONE, 1931) em sistema simbiótico em águas oligohalinas / Valdemir Queiroz de Oliveira. - 2021.
58 f.

Orientador: Alfredo Olivera Galvez.

Coorientador: Luis Otavio Brito da Silva.

Inclui referências e apêndice(s).

Dissertação (Mestrado) - Universidade Federal Rural de Pernambuco, Programa de Pós-Graduação em Recursos Pesqueiros e Aquicultura, Recife, 2021.

1. Equilíbrio iônico. 2. água oligohalina. 3. desempenho zootécnico. 4. minerais. 5. farelo de arroz fermentado. I. Galvez, Alfredo Olivera, orient. II. Silva, Luis Otavio Brito da, coorient. III. Título

CDD 639.3

UNIVERSIDADE FEDERAL RURAL DE PERNAMBUCO
PRÓ-REITORIA DE PESQUISA E PÓS-GRADUAÇÃO
PROGRAMA DE PÓS-GRADUAÇÃO EM RECURSOS PESQUEIROS E
AQUICULTURA

EFEITO DA FREQUÊNCIA DO AJUSTE EQUILÍBRIO IÔNICO NO
DESEMPENHO ZOOTÉCNICO DO *Litopenaeus vannamei* (BOONE, 1931) EM
SISTEMA SIMBIÓTICO EM ÁGUAS OLIGOHALINAS

Valdemir Queiroz de Oliveira

Dissertação julgada adequada para
obtenção do título de mestre em
Recursos Pesqueiros e Aquicultura.
Defendida e aprovada em 08/12/2021
pela seguinte Banca Examinadora.

Prof. Dr. Alfredo Olivera Gálvez

Orientador

Departamento de Pesca e Aquicultura - Universidade Federal Rural de Pernambuco

Prof^a. Dr^a. Juliana Ferreira dos Santos

Membro Interno

Departamento de Pesca e Aquicultura - Universidade Federal Rural de Pernambuco

Prof^a. Dr^a. Gelcirene de Albuquerque Costa

Membro Externo

Departamento de Pesca e Aquicultura - Universidade Federal Rural de Pernambuco

Agradecimentos

Gostaria de agradecer à Deus por me permitir esta oportunidade de novos conhecimentos profissionais.

À minha família, Zete e Arthur, por estarem sempre presentes durante esta jornada, e aos meus pais, por permitirem que seus filhos sempre estudassem, e sempre incentivando a estudar mais.

Ao meu orientador, professor Alfredo Olivera, e minha co-orientadora, Alitieni Pereira, por permitir a realização deste trabalho e transmitir ensinamentos e apoio durante esta jornada.

Ao professor Luis Otávio, um amigo e companheiro com quem aprendi um pouco sobre a carcinicultura e pelas conversas e troca de informações que me permitiu concluir esta etapa da minha vida.

À minha segunda família que tive o prazer de conhecer durante esta jornada, Otávio Augusto, Renata Farias, Elizabeth Santos, Marcele Trajano, Eduarda Almeida e Genison, pelas conversas, incentivo e ajuda, sem vocês este sonho seria bem mais difícil.

À equipe do Laboratório de carcinicultura (LACAR/UFRPE) pela ajuda no manejo durante o experimento.

À Embrapa Meio-Norte, por permitir aumentar este conhecimento profissional e que eu possa devolvê-lo à sociedade.

Aos membros da banca, prof. Prof^a. Dr^a. Gelcirene de Albuquerque Costa, prof.^a Dr.^a Suzianny Cabral, prof.^a Dr.^a Juliana Ferreira e prof.^a Dr.^a. Jéssika Lima de Abreu, pelas contribuições para o amadurecimento da dissertação.

A todos que de alguma maneira contribuíram para a realização deste trabalho, meu muito obrigado!

RESUMO

O cultivo do camarão marinho *Litopenaeus vannamei* em água de baixa salinidade é uma prática em crescente desenvolvimento no mundo e Brasil. Essa prática é possível devido as altas concentrações de dureza total e alcalinidade total das fontes de água, além de alguns ajustes realizados nos principais íons necessários para o desenvolvimento do animal. Estudos relatam que esses íons são utilizados pelos camarões para a realização de reações metabólicas, ao processo de ecdise, sendo absorvidos para a mineralização do exoesqueleto do animal, papel importante na ativação da enzima Na^+/K^+ -ATPase, responsável pelo transporte ativo de íons entre os meios extra e intracelulares entre outras. Entretanto não se sabe a frequência necessária que estes íons devem ser repostos no ambiente de cultivo, diante disso, o presente estudo visou avaliar o efeito de diferentes frequências de ajuste iônico no cultivo de camarão marinho em baixa salinidade, assim como verificar o efeito desta correção na composição mineral do floco microbiano e da composição corporal do camarão marinho *L. vannamei* cultivado em sistema simbiótico. Foi realizado um cultivo experimental ao longo de 40 dias, onde foram estocadas pós larvas a uma densidade de 2000 cam.m⁻³, em 5 tratamentos: SW (água do mar- salinidade 31), SWD (água do mar diluída- salinidade 2.3), 1IA (salinidade 2.3- ajuste de íons para atingir a relação 1:3:1 Ca:Mg:K apenas no 1º dia) – 2IA (salinidade 2.3- ajuste de íons para atingir a relação 1:3:1 Ca:Mg:K no 1º e 20º dia) e 3IA- (salinidade 2.3- ajuste de íons para atingir a relação 1:3:1 Ca:Mg:K no 1º, 10º e 20º dia). Os camarões foram alimentados na frequência de 4x ao dia com ração de 45% de proteína bruta e realizou-se fertilização orgânica com farelo de arroz (20g m⁻³ < 300 micra), açúcar demerara (2 g m⁻³), bicarbonato de sódio (4 g m⁻³) e 0.5 g m⁻³ de mix de bactérias contendo: *Bacillus subtilis*, *B. licheniformis*, *Saccharomyces* sp. e *Pseudomonas* sp. in a total of 5.5 to 6.5 × 10⁷ CFU g⁻¹, sendo essa mistura foi submetida por 24 horas ao processo anaeróbico e em seguida ao aeróbico (24 horas), realizadas a cada 3 dias. Os ajustes dos íons para manter a relação 1:3:1 Ca:Mg:K foram realizados utilizando cloreto de magnésio (10,5 % Mg), cloreto de potássio (51,4% K) e carbonato de cálcio (30,4%Ca). As variáveis de qualidade de água (NAT, NO₂-N, NO₃-N, PO₄, alcalinidade total, dureza total, cálcio, magnésio, potássio, sódio, sulfato, cloreto) e as biometrias foram mensuradas a cada 10 dias. A composição mineral dos flocos e do camarão foram mensuradas ao final dos 40 dias de cultivo. Os compostos nitrogenados se mantiveram dentro dos níveis recomendados (0,66 mg/L NAT, 0,52 mg/L NO₂-N, 1,91 mg/L NO₃-N) para o cultivo do *L. vannamei* em água

oligohalinas em sistema simbiótico. Os parâmetros de desempenho zootécnico (1,14 a 1,18 g de peso médio; 88,61 a 94,44 % de sobrevivência; 2,10 a 2,15 kg/m³ de produtividade) entre os tratamentos de baixa salinidade não diferiram significativamente quando comparados entre si, entretanto os resultados foram superiores aos encontrados para o tratamento de alta salinidade (1,02 g de peso médio; 84,72 % de sobrevivência; 1,73 kg/m³ de produtividade). Ao avaliar a concentração de minerais no floco microbiano e na biomassa do camarão foi observado que a frequência da realização do ajuste iônico também não influenciou significativamente essas concentrações, diferindo apenas quando comparada a composição dos elementos da água marinha. Ao final do cultivo, os camarões foram submetidos a um teste de estresse amoniacal durante 96h para avaliar a resistência dos juvenis, obtendo 100% de sobrevivência após o período de prova em todos os tratamentos. Dessa forma, com base nos resultados obtidos, pode-se produzir juvenis de *L. vannamei* em água do mar diluída (salinidade 2.3; 38,0 mg Ca/L, 96,0 mg Mg/L, 48,3 mg K/L, 92,33 mg Alcalinidade total/L e 490,13 mg dureza total/L, relação Ca:Mg:K 1:2,5:1,3) sem a realização de ajustes iônicos e sem troca de água ao longo de 40 dias na fase de berçário em sistema de simbiótico.

PALAVRAS-CHAVE: Equilíbrio iônico; água oligohalina; desempenho zootécnico; minerais; farelo de arroz fermentado.

ABSTRACT

The cultivation of the marine shrimp *Litopenaeus vannamei* in low salinity water is an increasingly developing practice. This practice is possible due to the adjustments made to the main ions necessary for the animal's development. Studies report that these ions are used by shrimp to carry out metabolic reactions, as well as in the biofloc system, they stimulate the stabilization of the system and the formation of microbial flakes. However, the frequency that these ions must be replaced in the system is not known. Therefore, the present study aimed to evaluate the effect of different frequencies of ionic adjustment in the cultivation of marine shrimp in low salinity, as well as to verify the effect of this correction on the composition mineral of the biofloc and body composition of marine shrimp *L. vannamei*. An experimental cultivation was carried out over 40 days, the larvae were stored after larvae at a density of 2000cam.m⁻³, in 5 treatments: SW (sea water-salinity 31), SWD (diluted sea water - salinity 2.3), 1IA (salinity 2.3 - adjustment only on the 1st day) - 2IA (salinity 2.3 - adjustment on the 1st and 20th day) and 3IA - (salinity 2.3 - adjustment on the 1st, 10th and 20th day). The water quality variables remained within the recommended levels for the cultivation of *L. vannamei* in a biofloc system. The zootechnical performance parameters between the low salinity treatments did not differ significantly when compared to each other, however the results were superior to those found for the high salinity treatment. When evaluating the concentration of minerals in the biofloc and biomass of shrimp, it was observed that the frequency of the ionic adjustment did not significantly influence these parameters either, differing only when compared to the composition of elements in seawater. At the end of culture, the animals were subjected to an ammonia stress test for 96h, however, all treatments obtained 100% survival after the test period. Thus, based on the results obtained, the use of seawater diluted to salinity 2.3 without performing ionic adjustments throughout the cultivation of *L. vannamei* in the nursery phase does not represent losses to zootechnical indices, and it is possible to obtain the same production rate and cost reduction related to fertilizers used to carry out ionic corrections.

KEYWORDS: Ionic equilibrium; oligohaline water; zootechnical performance; minerals; rice bran fermented.

Sumário

RESUMO	4
ABSTRACT	6
1. INTRODUÇÃO	8
1.2 OBJETIVOS	10
1.2.1 Objetivo geral:	10
1.2.2 Objetivos específicos:	11
1.2. Hipóteses	11
2. ARTIGO CIENTÍFICO - Effect of ionic adjustment frequency in low salinity water on zootechnical performance, water quality and mineral composition of <i>Litopenaeus vannamei</i> in a synbiotic nursery system	12
Abstract	14
2.1. Introduction	16
2.2. Material and methods	18
<i>Experimental design</i>	18
<i>Experimental conditions</i>	19
<i>Shrimp stocking, feeding and monitoring</i>	20
<i>Ionic adjustment</i>	21
<i>Water quality</i>	22
<i>Stress tests</i>	23
<i>Mineral composition (Floc and Shrimp)</i>	23
<i>Statistical analysis</i>	24
2.3. Results	25
<i>Shrimp zootechnical performance</i>	25
<i>Water quality</i>	25
<i>Ions</i>	26
<i>Microbial floc mineral composition</i>	26
<i>Mineral in shrimp biomass</i>	27
<i>Stress test</i>	27
2.4. Discussion	27
2.5. Conclusion	31
Acknowledgements	31
Funding sources	32
2.6. References	33
3. REFERENCIA BIBLIOGRÁFICA	52

1. INTRODUÇÃO

1.1 – CONTEXTUALIZAÇÃO DA PESQUISA

A produção de camarão marinho afastado da costa utilizando águas oligohalinas (salinidade 0,5 a 3,0 g L⁻¹) e mesohalinas (salinidade 3,0 a 16,5 g L⁻¹) com origem no lençol freático, reservatórios artificiais, lagos e rios ou salinizada artificialmente tem aumentado significativamente no mundo e no Brasil, explorando novas fronteiras do agronegócio (Brito et al, 2019). Diversos autores demonstram viáveis a produção de camarão marinho longe da costa, como: Samocha et al. (1998) (EUA), Van Wyk et al. (1999) (EUA), Nunes e López (2001) (Brasil e Equador), Boyd, 2001 (Tailândia, EUA, Equador), Samocha et al. 2001 (EUA e Israel), Boyd et al. (2002) (Tailândia, Equador, Brasil e EUA), Boyd e Thunjai (2003) (China, Tailândia, Equador e EUA), McNevin et al. (2004) (EUA), entre outros. Em 2018 a produção em áreas interiores já representou aproximadamente 39% (3.653 milhões de toneladas) da produção mundial de crustáceos (FAO, 2020). No Brasil, esta propensão também pode ser observada em estados como Alagoas, Ceará, Paraíba e Sergipe tem aumentado o número de produtores de camarões em empreendimentos distante do mar (IBGE, 2020). Esses resultados em água interiores deve ser a fácil adaptação dos camarões marinhos às condições de diversas salinidades, temperaturas e altas concentrações de compostos nitrogenados (Ponce-Palafox et al., 1997; Laramore et al., 2001; Lin e Chen, 2003).

Apesar de promissores resultados zootécnicos e econômicos, os corpos hídricos naturais interiores podem apresentar grandes variações de íons em sua composição, sendo influenciados por fatores geológicos e climáticos (Boyd e Thunjai, 2003; Boyd, 2015). A água do mar é composta principalmente pelos íons cloreto, sódio, sulfato, magnésio, cálcio, potássio e bicarbonato, o que corresponde à 99,75% da concentração dos minerais presentes, enquanto e outros 46 minerais correspondem à 0,25% (Hem, 1967). Entretanto, em águas interiores ocorrem modificações na concentração desses sais (Boyd, 2007).

A baixa disponibilidade de alguns íons para animais proporcionam uma maior gasto energético para manter as concentrações de sais nas células através da pressão osmótica (Schmidt-Nielsen, 1990; Boyd, 2015), além disso, vários íons são indispensáveis para fisiologia dos camarões *Litopenaeus vannamei*, como cálcio e magnésio para o processo de ecdise (Boyd e Tucker, 1998), sódio e potássio para o crescimento, osmorregulação e imunidade (Liu et al., 2016; Li et al., 2017), podendo em

baixa disponibilidade desses íons proporcionar elevadas taxas de mortalidade e limitações no crescimento dos camarões (Saoud et al., 2003; Roy e Davis, 2010).

Gao et al. (2016) relacionou a diminuição de salinidade com um menor peso médio final, imunodepressão e a redução da quimiotripsina, inibindo a taxa de síntese digestiva em juvenis de camarões. Essa redução de salinidade da água, proporciona um estresse salino que aumenta os níveis transcritos das enzimas osmorregulatórias Na^+ K^+ ATPase e anidrase carbônica (CA) em brânquias de crustáceos (Gao et al., 2012; 2016).

São necessárias concentração mínima dos íons Na^+ , Cl^- , K^+ , Ca^{2+} , Mg^{2+} , SO_4^{2-} e HCO_3^- e proporção entre eles relativas à água do mar, sendo necessário realizar o ajuste nas proporções entre os íons: Na^+ : Mg^{+2} : Ca^{+2} : K^+ de 27:3:1:1, garantindo assim condições para o crescimento e a sobrevivência do *L. vannamei* em sistemas com baixa salinidade (Boyd e Thunjai, 2003; Boyd, 2018). Essas concentrações podem ser ajustadas através da biocompensação mineral, na qual os sais são incorporados a ração do animal ou por meio da compensação na água do sistema, sendo utilizados fertilizantes minerais (Cheng et al., 2006; Liu et al., 2016).

A necessidade de realizar as correções de acordo com o perfil iônico da água podem variar as concentrações de fertilizantes necessárias para realização de tal ajuste. Aumentando assim a importância da determinação da frequência em que esses fertilizantes devem ser reaplicados, pois a frequência de compensação iônica artificialmente irá influenciar os custos de produção e pode tornar inviável a atividade dependendo do preço de comercialização do produto final (Galkanda-Arachchige et al., 2020; Pinto et al., 2020), entretanto esse custo de produção pode ser diluído durante vários ciclos de cultivo, quando utiliza-se os sistemas de mínima troca de água ou reduzidos quando utilizamos berçários, como o objetivo de produzir juvenis mais aclimatadas as baixas concentrações de íons e com melhor status imune.

Porém a utilização de sistema caracterizados pela mínima troca de água podem acarretar o aumento na concentração de compostos nitrogenados (amônia e nitrito) (Burford et al. 2003) provenientes da mineralização do resíduo alimentar e da excreção dos animais (Viadero Jr et al., 2005), principalmente nas primeiras semanas cultivo, quando as bactérias heterotróficas e nitrificantes ainda não estão bem estabelecidas. Elevadas concentrações de compostos nitrogenados são tóxicas para animais aquáticos (Ray et al., 2011), e a utilização de sistema com baixa salinidade pode então aumentar o

efeito da toxicidade da amônia e nitrito influenciando negativamente no crescimento e na sobrevivência do *L. vannamei* (Lin e Chen, 2001; 2003).

Para redução dos compostos nitrogenados, o sistema simbiótico é importante ferramenta para formação dos flocos microbianos, resultantes da fermentação e respiração microbiana de farelos vegetais por bactérias (*Bacillus*, *Bifidobacterium* e *Lactobacillus*) e/ou leveduras (*Saccharomyces cerevisiae*) (Dos Santos et al., 2022; Pimentel et al., 2022), produzindo ácidos orgânicos, vitaminas do grupo do complexo B ou antibióticos naturais, e melhorando a solubilidade das fontes de carboidratos (monossacarídeos, dissacarídeos e polissacarídeos) (Asgharian et al., 2016).

A utilização dos processos anaeróbicos (fermentação) e/ou aeróbios (respiração) com microorganismos (*Lactobacillus*, *Pediococcus*, *Enterococcus*, *Bacillus*, *Saccharomyces*, entre outros) está entre as formas de aumentar a disponibilidade de nutrientes de uma matéria-prima, especialmente os cereais. Durante a fermentação, há uma redução do teor de fibra e um aumento da biomassa microbiana. Esses microorganismos transformam os constituintes químicos das matérias-primas através da quebra de moléculas orgânicas complexas, como polissacarídeos, em moléculas mais simples, aumentando assim a biodisponibilidade e solubilidade dos nutrientes, tornando-os mais nutritivos (Azevedo et al., 2015) e permitindo a formação de ácidos orgânicos, reduzindo o pH e alterando o equilíbrio da microbiota intestinal reduzindo assim o crescimento de microorganismos patogênicos (Brito et al., 2013; Flesch et al., 2014).

Diante disto, o objetivo deste trabalho foi avaliar o efeito da frequência do ajuste iônico em águas oligohalinas na fase de berçários sobre o desempenho zootécnico, qualidade de água, composição mineral dos flocos microbianos e do camarão *L. vannamei* em sistema de simbiótico.

1.2 OBJETIVOS

1.2.1 Objetivo geral:

Determinar a melhor frequência de ajuste do equilíbrio iônico no desempenho zootécnico, do *Litopenaeus vannamei* em águas oligohalinas utilizando sistema simbiótico.

1.2.2 Objetivos específicos:

- Avaliar o desempenho zootécnico das pós-larvas de *L vannamei* submetidos à diferentes frequências do ajuste no equilíbrio iônico em sistemas simbióticos;
- Avaliar a frequência de ajuste do equilíbrio iônico na qualidade de água durante a fase de berçário de *L vannamei* em sistemas simbióticos;
- Avaliar a resistência das pós-larvas de *L vannamei*, cultivadas com diferentes frequências do ajuste no equilíbrio iônico, ao teste de estresse osmótico e teste de estresse amoniacal ao final da fase de berçário em sistema simbiótico;
- Avaliar o efeito da frequência do ajuste no equilíbrio iônico na composição mineral do floco microbiano em sistemas simbióticos;
- Avaliar o efeito da frequência do ajuste no equilíbrio iônico na composição mineral do *L vannamei* em sistemas simbióticos.

1.2. Hipóteses

- A frequência do ajuste iônico nas relações entre os cátions Ca:Mg:K na água oligohalina proporcionará melhor desempenho zootécnico dos camarões marinhos;
- A frequência do ajuste iônico nas relações entre os cátions Ca:Mg:K na água oligohalina proporcionará modificação nos compostos nitrogenados da água.

2. ARTIGO CIENTÍFICO - Effect of ionic adjustment frequency in low salinity water on zootechnical performance, water quality and mineral composition of *Litopenaeus vannamei* in a synbiotic nursery system

Artigo científico a ser submetido a revista Aquaculture

ISSN: 0044-8486

Todas as normas de redação e citação deste artigo, atendem aquelas estabelecidas pela referida revista.

Valdemir Queiroz de Oliveira^a; Otávio Augusto Lacerda Ferreira Pimentel^b; Caio Rubens do Rêgo Oliveira^c; Elizabeth Pereira dos Santos^c; Alitienne Moura Lemos Pereira^d; Alfredo Olivera Gálvez^c; Luis Otavio Brito^{c*}

^a Embrapa Meio-Norte, Teresina, Brazil. valdemir.queiroz@embrapa.br

^b Estação Marinha de Aquicultura, Instituto de Oceanografia, Universidade Federal do Rio Grande, Rio Grande, Brazil. otavio.pimentel@yahoo.com

^c Departamento de Pesca e Aquicultura, Universidade Federal Rural de Pernambuco,, Recife, Brazil. caiorubens90@nevada.unr.edu; santos.e.p.26@gmail.com; alfredo_oliv@yahoo.com; luis.obsilva@ufrpe.br

^d Embrapa Tabuleiros Costeiros, Aracaju, Brazil. alitienne.pereira@embrapa.br

*Corresponding author: Luis Otavio Brito, Departamento de Pesca e Aquicultura, Universidade Federal Rural de Pernambuco, Dois Irmãos, Recife, Pernambuco 52171-900, Brazil. Tel: +55 81 9990-2553, *e-mail address*: luis.obsilva@ufrpe.br

Abstract

Ions are important for metabolic processes in marine shrimps, but it is not known the need to maintain levels without affecting shrimp growth. The purpose of this work was to evaluate the effect of ionic adjustment frequencies in low salinity water in a synbiotic nursery system as well as verify its effects on the microbial flocs and *Litopenaeus vannamei* mineral composition. The experiment was carried out for 40 days, with a density of 2000 post-larvae m⁻³. Thus, following treatments were tested, all in triplicate: SW (seawater - salinity 31 g L⁻¹); SWD (seawater diluted to a salinity of 2.3 g L⁻¹); 1IA (salinity 2.3 g L⁻¹- ionic adjustment to achieve Ca:Mg:K ratio of 1:3:1 only on the 1st day); 2IA (salinity 2.3 g L⁻¹ - ionic adjustment to achieve Ca:Mg:K ratio of 1:3:1 on the 1st and 20th day) and 3IA - (salinity 2.3 g L⁻¹ - ionic adjustment to achieve Ca:Mg:K ratio of 1:3:1 on the 1st, 10th and 20th day). Shrimps were fed four times a day with a 45% protein diet. The synbiotic fertilization protocol used a mix of *Bacillus subtilis*, *B. licheniformis*, *Saccharomyces* sp. and *Pseudomonas* sp. in a total of 5.5 to 6.5 × 10⁷ CFU g⁻¹ every 3 days. The nitrogen compounds remained within the recommended levels (NAT = 0.66 mg L⁻¹, NO₂⁻-N = 0.52 mg L⁻¹, NO₃⁻-N = 1.91 mg L⁻¹) for *L. vannamei* reared in low salinity water. Mean weight survival and yield (mean weight: 1.14 to 1.18 g; survival: 88.61 to 94.44 % and yield: 2.10 to 2.15 Kg m⁻³) were not significantly different in low salinity treatments but were higher than those found in SW treatment (mean weight = 1.02 g; survival = 84.72%; and yield = 1.73 Kg m⁻³). Mineral concentration in microbial floc and shrimp biomass did not differ among the frequencies tested, only differing when compared to SW treatment. To evaluate the juvenile's resistance, an ammoniacal stress test was performed during 96h, obtaining 100% survival for all treatments. According to the results, *L. vannamei* juveniles can be produced in diluted seawater (salinity 2.3; Ca²⁺ = 38.0 mg L⁻¹, Mg²⁺ = 96.0 mg L, K⁺ = 48.3 mg L⁻¹,

total alkalinity = 92.33 mg L⁻¹ and total hardness = 490.13 mg L⁻¹, Ca:Mg:K ratio 1:2.5:1.3) without ionic adjustments and without water exchange for 40 days in the nursery phase using synbiotic system without harming production parameters.

Keywords: growth; stress test; Ca:Mg:K; mineral; nitrogen.

2.1.Introduction

Although the production of *Litopenaeus vannamei* is predominantly carried out in a marine or estuarine environment, there are already several productive results in places far from the sea, such as the Gobi desert in China, considered the furthest place from the sea on earth (FAO, 2020). In 2018, the marine shrimp production in inland areas corresponded to 3.653 million tons (52% of the world production of crustaceans) (FAO, 2020). These farms in areas with such different environmental characteristics are only feasible because *L. vannamei* is an easily adaptable species, supporting an extensive temperature and salinity gradient, as well as high concentrations of nitrogen compounds (Ponce-Palafox et al., 1997; Laramore et al., 2001; Lin and Chen, 2003).

The ease and reduced cost of inland areas are the main motivations for shrimp farming implementation in such lands, when compared to coastal lands, those inland areas present greater flexibility for acquisition of environmental licenses as well as access to reduced energy taxes.

These indoor crops are carried out using oligohaline (0.5 to 3.0 g L⁻¹ salinity) and mesohaline (3.0 to 16.5 g L⁻¹ salinity) waters from groundwater, artificial reservoirs, lakes and rivers, or artificially salt flats appear as a strategy of using inadequate water resources for agriculture and human consumption, increasing significantly the agribusiness borders.

Despite promising zootechnical and economic results, inland natural water bodies can present great variations in their composition, and this water profile is influenced by geological and climatic factors. (Boyd and Thunjai, 2003; Boyd, 2015).

The low availability of some ions for animals causes a greater energy expenditure to equalize the concentration of salts inside the cell through osmotic pressure (Schmidt-Nielsen, 1990; Boyd, 2015), in addition, several ions are associated with metabolic and chemical reactions. Physiological characteristics of *L. vannamei*, such as: calcium (Ca²⁺)

and magnesium (Mg^{2+}) participate in the ecdysis process (Boyd and Tucker, 1998), sodium (Na^+) and potassium (K^+) act on growth, osmoregulation and immunity (Liu et al., 2016; Li et al., 2017). When the *L. vannamei* culture is carried out in environments that present a low concentration of these ions, it can cause high mortality rates and limitations in the growth of shrimp. (Saoud et al., 2003; Roy and Davis, 2010).

Minimum concentrations of Na^+ , chloride (Cl^-), K^+ , Ca^{2+} , Mg^{2+} , sulfate (SO_4^{2-}) and bicarbonate (HCO_3^-) ions and a proportion among them relative to seawater are required, and it is necessary to adjust the proportions among the ions: $Na^+ : Mg^{2+} : Ca^{2+} : K^+$ of 27:3:1:1, thus ensuring conditions for the *L. vannamei* growth and survival in systems with low salinity (Boyd and Thunjai, 2003; Boyd, 2020). These concentrations can be adjusted through mineral bio compensation, a process in which the salts are incorporated into the animals feed or into the system's water, using mineral fertilizers (Cheng et al., 2006; Liu et al., 2016).

The concentration of fertilizers required to make the corrections are determined according to the ionic profile of the water. Thus, increasing the importance of determining the frequency at which these fertilizers should be reapplied, as the frequency of ionic compensation will artificially influence production costs and may make the activity unfeasible depending on the marketing price of the final product. (Galkanda-Arachchige et al., 2020; Pinto et al., 2020). However, this production cost can be diluted during several farming cycles, using systems of minimum water exchange, for this, a good control of nitrogenous compounds in low salinity is necessary, due to the toxicity of these compounds to cultivated marine shrimps.

Synbiotic system, on the other hand, results from the decomposition of products of plant origin, which are mostly wheat, soybean or rice bran, by bacteria (*Bacillus*, *Bifidobacterium* and *Lactobacillus*) and/or yeasts (*Saccharomyces cerevisiae*), into a

controlled aerobic and/or anaerobic process, which produces organic acids, B-complex vitamins or natural antibiotics, improving the solubility of carbohydrate sources (monosaccharides, disaccharides and polysaccharides). These microorganisms transform the chemical constituents of raw materials by breaking complex organic molecules, such as polysaccharides, into simpler molecules, thus increasing the bioavailability and solubility of nutrients, making them more nutritious. Consequently, they increase the control of nitrogenous compounds by increasing the microbial biomass in the system (heterotrophic and nitrifying bacteria).

Therefore, the aim of this study was to evaluate the effect of the ionic adjustment frequency in oligohaline water on the zootechnical performance, water quality, *L. vannamei* mineral and microbial floc composition in a synbiotic nursery system.

2.2. Material and methods

Experimental design

This study was carried out for 40 days at the Shrimp Culture Laboratory (LACAR), of the Fisheries and Aquaculture Department of the Rural Federal University of Pernambuco (UFRPE), Brazil. The experimental design was completely randomized with five treatments: SW (seawater - salinity 31 g L⁻¹); SWD (seawater diluted to a salinity of 2.3 g L⁻¹); 1IA (salinity 2.3 g L⁻¹- ionic adjustment to achieve Ca:Mg:K ratio of 1:3:1 only on the 1st day); 2IA (salinity 2.3 g L⁻¹ - ionic adjustment to achieve Ca:Mg:K ratio of 1:3:1 on the 1st and 20th day) and 3IA- (salinity 2.3 g L⁻¹ - ionic adjustment to achieve Ca:Mg:K ratio of 1:3:1 on the 1st, 10th and 20th day), all in triplicate.

Experimental conditions

The treatments 1IA, 2IA, 3IA e SWD used seawater with 35 g L⁻¹ diluted with freshwater 0.16 g L⁻¹ to a salinity of 2.3 g L⁻¹. The water inside the experimental units was prepared 15 days prior to animal stocking. This procedure was carried out by filling 15% (9 L) of the microbial floc tanks (TAN = 0.49 mg L⁻¹, N-NO₂⁻ = 0.70 mg L⁻¹, N-NO₃⁻ = 0.81 mg L⁻¹, total hardness = 763 mg CaCO₃ L⁻¹, total alkalinity = 88 mg CaCO₃ L⁻¹), and 85% (51 L) of water treated with sodium hypochlorite in a concentration of 15 g m⁻³ of activated chloride. Then, the synbiotic fertilization was applied. The fertilizer was submitted to an anaerobic process (24 h) and an aerobic phase (24 h) which was composed of 20 g m⁻³ of rice bran (< 200 µm), 2 g m⁻³ of sugar, 4 g m⁻³ of sodium bicarbonate and 0.5 g m⁻³ of commercial bacterial mix [5.5 to 6.5 × 10⁷ colony-forming units (CFU) g⁻¹, containing: *Bacillus subtilis*, *Bacillus licheniformis*, *Saccharomyces* sp. e *Pseudomonas* sp. (Kayros Agrícola and Ambiental, Brazil) and chlorinated water [20 mg L⁻¹ active chlorine (60%), then unchlorinated through aeration] in a proportion to 10 × the amount of rice bran. Totaling seven applications, with a two-day interval among fertilizations. The organic fertilizer was added to the experimental treatments every three days during culture until settleable solids reached 5 ml L⁻¹.

The experimental units (60 L), were constantly aerated (dissolved oxygen > 5.0 mg L⁻¹), temperature maintained at (~31°C; Hopar Sh-608 heater 100 W), 12:12-h photoperiod and mean luminance of 8.65 µmol photons m⁻² s (Equitherm Lux-204). No water exchange was performed during the experiment, and dechlorinated freshwater was added to replace evaporation loss four times a week. The artificial substrate was composed of mollusk shells (*Anomalocardia brasiliiana*), which covered ≈ 28.12% of the bottom area (25 × 24 × 5 cm, width × height × depth) and corresponded to ≈ 3.36% of the experimental unit's useful volume.

Shrimp stocking, feeding and monitoring

The post larvae (PL10 1.18 ± 0.4 mg) were acquired from a commercial hatchery (Aquatec, Rio Grande do Norte, Brazil), which was produced in water with salinity at 35 g L⁻¹. Prior to experiment, the PL10 were acclimated to salinity condition of 2.5 g L⁻¹ for 10 days. At the end of the acclimatization period, the PL 20 (10 ± 3 mg) were randomly selected and stocked at a density of 2000 PL's m⁻³ in experimental units.

The post larvae were fed with commercial feed of 45% crude protein, 7% lipids, (0.3 - 0.6 mm, Inve Aquaculture Inc), during 1 day to 17days, and with 45% crude protein, 9.5% lipids, (0.8 – 1.3 mm, ADM Animal Nutrition Company), between 17 days to 40 days, four times a day (8:00 am, 11:00 am, 2:00 pm and 5:00 pm). Initially, a 27.2% daily feeding rate was adopted, which was gradually reduced until 4.8% of body weight during 40 days. The feeding rate was daily adjusted according to the shrimp feed consumption estimate and mortality rate in each experimental unit (Van Wyk et al., 1999).

Shrimp weight was monitored every 10 days to determine shrimp growth and adjust the amount of feed offered. At the end of the experimental time, biomass gain, mean final weight, specific growth rate (SGR), FCR, survival and yield were determined based on the following equations:

Biomass gain (g) = final biomass (g) – initial biomass (g);

Final weight (g) = final biomass (g)/number of individuals at the end of evaluation period;

SGR (% day⁻¹) = $100 \times [\ln \text{ final weight (g)} - \ln \text{ initial weight (g)}] / \text{time (days)}$;

FCR = feed supplied/biomass gain;

Survival (%) = (number of individuals at the end of evaluation period/ initial number of individuals) $\times 100$;

Yield (Kg m⁻³) = final biomass (Kg)/volume of experimental unit (m³).

Ionic adjustment

The ionic adjustment was performed in treatments (1IA, 2IA and 3IA), according to the result of water quality variables based on the Ca:Mg:K (1:3:1) ratios proposed by Boyd and Thunjai (2003). For the adjustment of these ions, commercial products were used: potassium chloride (KCl), calcium carbonate (CaCO₃) and magnesium chloride hexahydrate (MgCl₂ · 6H₂O). The addition of ions were made based on previous tests of the efficiency of the chemical compounds in increasing the concentration of ions in the water (applied in the experimental units in two applications: 50% in the morning and 50% in the afternoon). The test was performed in experimental units with 14 L of water with a salinity of 2.5 g L⁻¹, obtained from seawater diluted in freshwater, under constant aeration. The pH and ion concentration were analyzed before application and 72 h after application of the chemicals at a concentration of 100 g m⁻³ (Table 1).

Location of the table 1

The chemical compounds application on the experimental units to adjust the ionic concentration was performed according to the following equation (Samocho, 2019):

$$\text{Amount of product} = \frac{Fc - Ic}{TI} \times V$$

- Fc is the final concentration;
- Ic is the initial concentration;
- TI is the % increment of the target ion (decimal value);
- V is the tank volume (m³).

After analyzing the water in all treatments, ion concentrations in milliequivalent L^{-1} ($mEq L^{-1}$) were calculated to check cations and anions equilibrium. This calculation was made by the difference between the sum of the cations $mEq L^{-1}$ ($Na^{+} = 23 mg mEq^{-1}$; $K^{+} = 39.1 mg mEq^{-1}$; $Ca^{2+} = 20 mg mEq^{-1}$ and $Mg^{2+} = 12.15 mg mEq^{-1}$) and the sum of the anions $mEq L^{-1}$ ($HCO_3^{-} = 61 mg mEq^{-1}$; $Cl^{-} = 35.45 mg mEq^{-1}$ and $SO_4^{2-} = 48.03 mg mEq^{-1}$) for chemical equilibrium certification (Boyd, 2015). As a standard for certifying the accuracy of the analysis of these major ions, a balance error between cations and anions below 15% was adopted (Custodio and Llamas, 1983). This error was calculated using the following equation:

$$Error (\%) = \frac{|\Sigma cations - \Sigma anions|}{\Sigma cations + \Sigma anions} \times 200$$

Where:

- Σ cations: sum of cations (Na^{+} , K^{+} , Ca^{2+} e Mg^{2+});
- Σ anions: sum of anions (HCO_3^{-} , Cl^{-} e SO_4^{2-}).

Water quality

Dissolved oxygen (DO; $mg L^{-1}$) and temperature ($^{\circ}C$; YSI model 100, Yellow Springs, Ohio, USA), were monitored twice a day (at 08:00 AM and 04:00 PM). Salinity (salinity meter AZ, model 8371), pH (pH meter Akso, model A90), and settleable solids ($mL L^{-1}$; Imhoff cone) (Avnimelech, 2012) were monitored three times a week. Total ammonia nitrogen (TAN; APHA, 2012), nitrite-N (NO_2^{-} -N; Fries, 1971), nitrate-N (NO_3^{-} -N; APHA, 2012), orthophosphate (PO_4^{3-} ; APHA, 2012), total alkalinity (TA; APHA, 2012), total hardness (TH; APHA, 2012), Ca^{2+} (APHA, 2012), Mg^{2+} (APHA, 2012), Na^{+} (APHA, 2012), Cl^{-} (APHA, 2012), SO_4^{2-} (APHA, 2012) and K^{+} (Fries and Getrost, 1977) were monitored every ten days.

Stress tests

At the end of the experimental time, the shrimp juveniles were submitted to resistance tests for osmotic stress and water ammonia nitrogen concentration ($\text{NH}_3\text{-N}$). For the osmotic stress test, 10 shrimps were randomly collected in each experimental unit and were stocked in experimental units with 10 L of seawater (35 g L^{-1}), with aeration, for 30 min (SWD, 1IA, 2IA, 3IA) and with 10 L of freshwater (0 g L^{-1}), with aeration, for 30 min (SW). After this time, the animals were exposed to salinity of 2.5 g L^{-1} for 30 min (SWD, 1IA, 2IA, 3IA) and exposed to salinity of 35 g L^{-1} for 30 min (SW). Survival was then evaluated for each treatment (Burbano-Gallardo et al., 2015).

For the $\text{NH}_3\text{-N}$ resistance test, at the end of the experiment, 10 shrimps were randomly collected in each experimental unit and were transferred to experimental units containing 10 L of water salinity 2.5 g L^{-1} (SWD, 1IA, 2IA, 3IA), and 10 L of water salinity 35 g L^{-1} (SW) with $\text{NH}_3\text{-N}$ concentrations between 0.26 and 0.41 mg L^{-1} (Table 2), water temperature close to 28.2°C and pH close to 8.0. The $\text{NH}_3\text{-N}$ concentration was achieved by applying a stock solution of 10 g L^{-1} of NH_4Cl . The test was carried out for 96 h, and survival was measured every 24 h (Zhang et al., 2012).

*Location of the table 2**Mineral composition (Floc and Shrimp)*

At the end of experimental time the floc samples (30 g) were collected with a cylindrical mesh net of $50 \mu\text{m}$ for solids retention and shrimp whole body, in triplicate for mineral composition analyses. Prior to the processing of the analysis, the microbial flocs samples were dried in an oven until constant weight was obtained and then kept in a desiccator, while the whole shrimp samples were freeze-dried, then both were subjected

to wet digestion through nitro-perchloric mineralization and submitted to the respective readings. Potassium (K), calcium (Ca), magnesium (Mg), sodium (Na), copper (Cu), iron (Fe), manganese (Mn) and zinc (Zn), from atomic absorption spectrophotometry (Agilent Technologies, US - SpectrAA 55B), as for phosphorus (P) and sulfur (S), they were performed with a UV-VIS spectrophotometer (Thermo Scientific, FI – Genesys 10S UV-Vis), according to the methodology described in Nogueira and Souza (1998). The samples were analyzed in the Plant Nutrition Laboratory- Brazilian Agricultural Research Corporation (EMBRAPA).

Statistical analysis

The data were checked for homogeneity of variance with the Cochran test ($p < 0.05$) and normality using the Shapiro–Wilk test ($p < 0.05$). One-way variance analysis (ANOVA) was conducted to evaluate the zootechnical performance variables and proximal composition, and repeated measures ANOVA were used to compare water quality data, followed by the Duncan test ($p < 0.05$). For non-parametric statistical data, Kruskal–Wallis ($p < 0.05$) followed by Dunn's test (for the SGR data) and Friedman's test with Conover multiple comparison test with Holm–Bonferroni correction (for DO, SS, total alkalinity and NO_2^- -N data). Data analysis were performed using Statistica 12.5 software.

2.3. Results

Shrimp zootechnical performance

The zootechnical performance data at the end of the 40 days of experimental culture are summarized in Table 3. SW treatment (final weight = 1.02 g; survival = 84.72 %; yield = 1.7 Kg m⁻³ and SGR = 11.49 % day⁻¹) is lower as compared to low salinity treatments (SWD, 1IA, 2IA and 3IA). However, FCR showed no significant differences ($p > 0.05$) among treatments.

Location of the table 3

Water quality

There were significant differences ($p < 0.05$) in the water quality parameters (dissolved oxygen, SS, salinity, pH, total hardness, total alkalinity, TAN and NO₂⁻-N) among the five treatments (Table 4). The lowest levels of dissolved oxygen (were observed in the SW treatment (4.75 mg L⁻¹) when compared to the low salinity treatments (5.70 mg L⁻¹). The mean SS concentration was higher in the ionic adjustment treatments compared to SW and SDW, and the range was also greater in the low salinity with ionic adjustment. The pH was lower in 1IA, 2IA and 3IA when compared to SW, but they were similar between the low salinity treatments.

Total hardness and total alkalinity were higher in SW treatments compared to low salinity treatments. The total alkalinity in the ionic adjustment treatments did not show significant differences among treatments, but in the total hardness we observed an increase in the concentration with adjustment frequencies, as well as along the culture time. The orthophosphate values were higher than 30 mg L⁻¹, with higher concentrations in the ionic adjustment treatments.

The TAN and NO_2^- -N values in the low salinity treatments were lower than SW, but NO_3^- -N did not show significant differences among the treatments. The concentrations of TAN and NO_2^- -N were higher until the 20th day of culture, followed by decrease.

Location of the table 4

Ions

The ions Ca^{+2} (402.7 mg L⁻¹), K^+ (217.1 mg L⁻¹), SO_4^- (2,817.5 mg L⁻¹), Cl^- (19,970.2 mg L⁻¹) and Na^+ (11,035.5 mg L⁻¹) showed significant differences ($p < 0.05$) between the SW treatment and the other low salinity treatments (Ca^{+2} : 38 - 51,3 mg L⁻¹; Mg^{+2} : 96- 163,5 mg L⁻¹; K^+ : 48,3 – 61,5 mg L⁻¹), however, only the Na:K ratio was significantly different among treatments. The Mg:Ca ratio was similar among the SW treatment and the treatments with ions adjustment (1IA; 2IA and 3IA), however they were significantly different to the SWD. Among treatments in oligohaline water, only Mg^{+2} concentration was significantly different between treatments (Table 5).

Location of the table 5

Microbial floc mineral composition

Minerals in microbial floc at the end of the experimental time are summarized in Table 6. The levels of P (1.90 -4.25 g Kg⁻¹), Fe (621.65 - 1167.46 mg. kg⁻¹) and Mn (11.31 - 16.29 mg Kg⁻¹) did not differ significantly among treatments. The concentrations of K, Mg, Na, S, Cu and Zn were higher in the SW treatments as compared to the then low

salinity treatments, however the Ca concentration was similar among the SW and the treatments with ions adjustment of Ca:Mg:K.

Location of the table 6

Mineral in shrimp biomass

The data for the mineral in the shrimp biomass at the end of the experimental time are summarized in Table 7. The levels of P, K, Ca, Mg, Na, S, Cu, Fe, Mn and Zn were not significantly different in the low salinity groups. However, Mg was higher in the SW treatment when compared to SWD and 1IA. Fe was significantly lower in the SW treatment when compared to low salinity treatments.

Location of the table 7

Stress test

At the end (96 hours) osmotic and ammonia nitrogen concentration (NH₃-N). Stress test survival was 100% in all treatments.

2.4. Discussion

Dissolved oxygen, temperature and pH of the water remained, throughout the experimental period, as recommended for *Litopenaeus vannamei* culture in intensive systems with minimal water exchange (Samocha, 2019). The high total hardness concentration found in the SW treatment was expected (due to its high salinity and seawater composition) the difference found among the treatments came from the different ionic adjustment frequencies, since among the products used there was magnesium

chloride and calcium carbonate, thus causing an increase in Ca^{2+} and Mg^{2+} ions, which constitute the total hardness level (Boyd, 2020).

During the experimental time, alkalinity was maintained above $75 \text{ mg CaCO}_3 \text{ L}^{-1}$, as recommended by Boyd and Tucker (1998). Alkalinity is an important water quality parameter in marine shrimp culture, being related to the water buffering which causes a lower pH oscillation throughout the day (Boyd et al., 2016), as observed in Table 4. In addition, Alkalinity provides nutrients necessary for the development of chemoautotrophic bacteria which is responsible for cycling nitrogen compounds (Ebeling et al., 2006). In the present study, it is possible that the difference found in this variable among the SWD, 1IA, 2IA and 3IA treatments is related to the nitrifying bacteria communities established in the system, since the differences found in alkalinity reflect the differences observed in TAN concentration. In addition, the presence of nitrifying bacteria can be observed in all treatments due to NO_2^- -N and NO_3^- -N concentrations in the water as a result of the oxidation process carried out by these communities. The nitrogenous compounds concentrations were maintained as recommended for the marine shrimp culture, indicating a system stabilization (Ramirez-Rochin et al., 2017; Boyd, 2020). The efficient stabilization of the nitrogenous compounds was also reported by Pimentel et al. (2022), Santos et al., (2022) and Romano et al., (2018), who, when performing shrimp culture in low salinity water, attributed the efficiency of nitrogen cycling to the use of rice bran submitted to anaerobic (24h) and aerobic (24h) process.

The use of the synbiotic as an organic carbon source to maintain good water quality conditions by controlling the total ammonia nitrogen and settleable solids concentration contributes to maintaining the C:N ratio close to 6.5:1. This ratio is close to that found by Santos et al. (2022) who used the same fertilization strategy of the present study, where they associated the use of the synbiotic to the reduction of settleable solids

volume. The highest values of settleable solids found in this study were related to the treatments where ionic adjustment was performed. In addition, the shells used as an artificial substrate retained the solids, due to the water circulation between them, until dissociation, after solid saturation on the shell substrate.

Microbial flocs have their composition affected by several factors, such as the system in which they are produced and also by physical and chemical variables, such as salinity (Khanjani et al., 2020), since these particles also have ions in their composition (De Schryver et al., 2008). The use of different frequencies of ionic adjustment in the culture system does not influence the mineral composition of the major ions present in the microbial floc. In the present study, when evaluating the treatments in which the ion adjustment (1IA, 2IA and 3IA) were performed, significant differences were observed only in the levels of Mg^{2+} , with the composition of the other elements evaluated being similar. These minerals can be made available to animals through bioflocs, as these microbial flocs act as a supplemental food source for shrimp (Khanjani et al., 2020). Authors such as Cheng et al., (2005; 2006) observed benefits related to the adoption of mineral supplementation through the feed as a strategy to improve the zootechnical indexes in the culture of *L. vannamei* in low salinity water.

The marine shrimp *L. vannamei* culture in low salinity water has been put to the test, since studies show that the reduction in salinity affects production indices such as survival, average final weight, yield, among others (Esparza-Leal et al., 2016; Laramore et al., 2001). In general, this event is justified by the low salinity and, consequently, the low availability of the major ions for these animals, providing a greater energy expenditure to osmotic and ionic regulation, which can be reflected in the zootechnical performance indices throughout a culture (Roy et al., 2010). However, the different frequencies of ionic adjustment in salinity 2.3 g L^{-1} using a system with minimal water

exchange and synbiotic as organic carbon source do not significantly influence the productive indexes in the *L. vannamei* culture in the nursery phase, since zootechnical results as final mean weight, final biomass, yield and specific growth rate were superior to those found in the treatment using seawater (salinity 31 g L⁻¹).

The shrimp performance results of this study were superior to those found by Esparza-Leal et al., (2016) who evaluated the effect of salinity on the growth of *L. vannamei* and obtained a mortality rate of 100% in salinities 2 and 4 g L⁻¹, while Pimentel et al., (2022), who evaluated the effect of different ionic adjustment strategies on the *L. vannamei* growth performance in salinity 2.5 g L⁻¹, obtained average survival rates lower than those obtained in the present study, reporting an average rate between 82 and 87%. The results of final weight, FCR, SGR and yield reached in the present study exceed those found by the same author. Such differences may be associated with the amount of protein offered to the animals through the feed, since the present study used feed with greater availability of protein. In addition, the mean concentrations of the major ions (Ca⁺², Mg⁺² and K⁺) that were maintained in the culture reported by the author, are lower than the lowest mean concentrations found in the present study, and although it indicates that the minimum concentrations maintained in that study did not limit the growth of *L. vannamei* in low salinity conditions, it is known that these ions are related to the growth and survival of shrimp when they are in low salinity conditions (Roy et al. 2007; Zacarias, et al., 2018).

Regarding the mineral composition of the *L. vannamei* biomass, the different ionic adjustment frequencies did not influence this parameter. The amount of total minerals can be obtained by determining the amount of ash in a proximal analysis, comparing the composition of *L. vannamei* cultivated in marine water with the biomass of the same species cultivated in low salinity. Liang et al., (2008) did not observe the influence of salinity on the concentration of minerals in the muscle of the animals. It is possible that

the minerals absorbed or consumed by animals are used in metabolic and physiological reactions of the animal, since these ions found in water and feed play a relevant role in metabolic reactions, influencing growth and animal survival.

2.5. Conclusion

The ionic adjustment frequency of the Ca:Mg:K ratio to 1:3:1 did not result in zootechnical differences in the *L. vannamei* nursery using diluted seawater in a synbiotic system. In view of the results of the zootechnical index obtained in the present study, the use of marine water without performing ionic adjustment is equivalent to the results obtained with the performance of different frequencies of this adjustment. In this way, it is possible to carry out *L.vannamei* culture using the synbiotic based on rice bran, in a minimal water exchange system only in salinity 2.3 g L⁻¹, using diluted seawater, saving costs resulting from the ionic adjustments through inorganic fertilization.

Concentrations of Ca²⁺ = 38.0 mg L⁻¹, Mg²⁺ = 96.0 mg L⁻¹ and K⁺ = 48.3 mg L⁻¹ are sufficient for marine shrimp nurseries in low salinity using synbiotic system.

Although the ionic adjustment is performed in the water, the ratios between the ions found in the microbial flocs has other ratios different from those found in the water and needs to be better studied, as these minerals are ingested by shrimp and may supply some need for osmotic bio compensation.

Acknowledgements

The authors are grateful for the financial support provided by the Rural Federal University of Pernambuco (Pró-Reitoria de Pesquisa e Pós-Graduação - PRPPG 015/2018); Brazilian Agricultural Research Corporation/Embrapa (21188.005941/2018-04); Brazil's National Council for Scientific and Technological Development (CNPq) and

the Coordination for the Improvement of Higher Education Personnel (CAPES) for the scholarships granted (process numbers 88882.436206/2019-01; 8882.436207/2019-01; 88882.436231/2019-01) and CNPq also provided a grants to Alfredo Olivera Gálvez (PQ 308063/2019-8) and Luis Otavio Brito (PQ309669/2021-9).

Funding sources

This work was supported by Rural Federal University of Pernambuco (Pró-Reitoria de Pesquisa e Pós-Graduação - PRPPG 015/2018); Brazilian Agricultural Research Corporation/Embrapa (21188.005941/2018-04); Brazil's National Council for Scientific and Technological Development- CNPq (PQ 308063/2019-8; PQ309669/2021-9) and the Coordination for the Improvement of Higher Education Personnel (CAPES) for the scholarships granted (88882.436206/2019-01; 88882.436207/2019-01; 88882.436231/2019-01).

Credit author statement

Valdemir Queiroz de Oliveira: Investigation, Conceptualization, Methodology, Formal analysis, Writing - Original Draft. **Otávio Augusto Lacerda Ferreira Pimentel:** Investigation, Data Curation, Methodology, Writing - Review and Editing. **Caio Rubens do Rêgo Oliveira:** Data Curation, Resources. **Elizabeth Pereira dos Santos:** Formal analysis, Writing - Review and Editing. **Alitene Moura Lemos Pereira:** Resources, Writing - Review and Editing. **Alfredo Olivera Gálvez:** Methodology, Writing - Review and Editing. **Luis Otavio Brito:** Supervision, Methodology, Resources, Writing - Review and Editing.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

2.6. References

APHA, AWWA, Wef, 2012. Standard methods for the examination of water and wastewater, 22nd ed. American Public Health Association.

Avnimelech, Y., 2012. Biofloc technology – A practical guide book (2nd ed.). The World Aquaculture Society.

Boyd, C.E., 2020. Water quality: An introduction. Springer Nature, Cham, Switzerland.

Boyd, C.E., 2015. Water quality: An introduction, 2nd ed. Springer Nature.

Boyd, C.E., Thunjai, T., 2003. Concentrations of major ions in waters of inland shrimp farms in China, Ecuador, Thailand, and the United States. J. World Aquac. Soc. 34, 524–532. <https://doi.org/10.1111/j.1749-7345.2003.tb00092.x>

Boyd, C.E., Tucker, C.S., 1998. Pond Aquaculture Water Quality Management, 1998th ed. Chapman and Hall, London, England.

Boyd, C.E., Tucker, C.S., Somridhivej, B., 2016. Alkalinity and hardness: Critical but elusive concepts in aquaculture: Alkalinity and hardness. J. World Aquac. Soc. 47, 6–41. <https://doi.org/10.1111/jwas.12241>

Burbano-Gallardo, E., Imués-Figueroa, M.A., Gonzalez-Legarda, E.A., Brito, L.O.,

Galvez, A.O., Arana, L.A.V., 2015. Supervivencia de poslarvas de *Litopenaeus vannamei* sometidas a la prueba de estrés osmótico y su relación con el estado de muda. Revista de biología marina y oceanografía 50, 323–329.

Cheng, K.-M., Hu, C.-Q., Liu, Y.-N., Zheng, S.-X., Qi, X.-J., 2006. Effects of dietary calcium, phosphorus and calcium/phosphorus ratio on the growth and tissue mineralization of *Litopenaeus vannamei* reared in low-salinity water. Aquaculture 251, 472–483. <https://doi.org/10.1016/j.aquaculture.2005.06.022>

Cheng, K.-M., Hu, C.-Q., Liu, Y.-N., Zheng, S.-X., Qi, X.-J., 2005. Dietary magnesium requirement and physiological responses of marine shrimp *Litopenaeus vannamei* reared in low salinity water. Aquac. Nutr. 11, 385–393. <https://doi.org/10.1111/j.1365-2095.2005.00364.x>

Custodio, E., Llamas, M. R., 1983. Hidrología subterránea (2nd ed.). Ediciones Omega.

De Schryver, P., Crab, R., Defoirdt, T., Boon, N., Verstraete, W., 2008. The basics of bio-flocs technology: The added value for aquaculture. Aquaculture 277, 125–137. <https://doi.org/10.1016/j.aquaculture.2008.02.019>

Ebeling, J.M., Timmons, M.B., Bisogni, J.J., 2006. Engineering analysis of the stoichiometry of photoautotrophic, autotrophic, and heterotrophic removal of ammonia–nitrogen in aquaculture systems. Aquaculture 257, 346–358. <https://doi.org/10.1016/j.aquaculture.2006.03.019>

Esparza-Leal, H.M., Amaral Xavier, J.A., Wasielesky, W., Jr, 2016. Performance of *Litopenaeus vannamei* postlarvae reared in indoor nursery tanks under biofloc

conditions at different salinities and zero-water exchange. *Aquac. Int.* 24, 1435–1447. <https://doi.org/10.1007/s10499-016-0001-5>

FAO, 2020. The state of world fisheries and aquaculture 2020 (SOFIA): Sustainability in action. Food & Agriculture Organization of the United Nations (FAO), Rome, Italy.

Fries, J., 1971. Análisis de Trazas. Métodos fotométricos comprobados. E.Merck, 130p.

Fries, J., Getrost, H., 1977. Organic Reagents for Trace Analysis. E. Merck Darmstadt, 453p.

Galkanda-Arachchige, H.S.C., Roy, L.A., Davis, D.A., 2020. Evaluation of an alternative salt mixture to culture Pacific white shrimp (*Litopenaeus vannamei*) in inland aquaculture. *Aquac. Res.* 51, 3540–3550. <https://doi.org/10.1111/are.14691>

Khanjani, M.H., Sharifinia, M., Hajirezaee, S., 2020. Effects of different salinity levels on water quality, growth performance and body composition of Pacific white shrimp (*Litopenaeus vannamei* Boone, 1931) cultured in a zero water exchange heterotrophic system. *Ann. Anim. Sci.* 20, 1471–1486. <https://doi.org/10.2478/aoas-2020-0036>

Laramore, S., Laramore, C.R., Scarpa, J., 2001. Effect of low salinity on growth and survival of postlarvae and juvenile *Litopenaeus vannamei*. *J. World Aquac. Soc.* 32, 385–392. <https://doi.org/10.1111/j.1749-7345.2001.tb00464.x>

- Li, E., Wang, X., Chen, K., Xu, C., Qin, J.G., Chen, L., 2017. Physiological change and nutritional requirement of Pacific white shrimp *Litopenaeus vannamei* at low salinity. Rev. Aquac. 9, 57–75. <https://doi.org/10.1111/raq.12104>
- Liang, M., Wang, S., Wang, J., Chang, Q., Mai, K., 2008. Comparison of flavor components in shrimp *Litopenaeus vannamei* cultured in sea water and low salinity water. Fish. Sci. 74, 1173–1179. <https://doi.org/10.1111/j.1444-2906.2008.01637.x>
- Lin, Y.-C., Chen, J.-C., 2003. Acute toxicity of nitrite on *Litopenaeus vannamei* (Boone) juveniles at different salinity levels. Aquaculture 224, 193–201. [https://doi.org/10.1016/s0044-8486\(03\)00220-5](https://doi.org/10.1016/s0044-8486(03)00220-5)
- Liu, H., Tan, B., Yang, J., Chi, S., Dong, X., Yang, Q., 2016. Effects of aqueous Na/K and dietary K on growth and physiological characters of the Pacific white shrimp, *Litopenaeus vannamei*, reared in low-salt well water. Aquac. Res. 47, 540–553. <https://doi.org/10.1111/are.12513>
- Nogueira, A.R.A., Machado, P.L.O.A., Souza, C.A.F.S., Ferreira, J.R., 1998. Manual de laboratórios: solo, água, nutrição vegetal, nutrição animal e alimentos. EMBRAPA-CPPSE, 72p.
- Pimentel, O.A.L.F., Oliveira, V.Q., Oliveira, C.R. do R., Severi, W., Olivera Gálvez, A., Amado, A.M., Brito, L.O., 2022. Assessment of different ionic adjustment strategies in low-salinity water on the growth of *Litopenaeus vannamei* and microbial community stoichiometry in a synbiotic nursery system. Aquac. Res. 53, 50–62. <https://doi.org/10.1111/are.15552>

- Pinto, P.H.O., Rocha, J.L., do Vale Figueiredo, J.P., Carneiro, R.F.S., Damian, C., de Oliveira, L., Seiffert, W.Q., 2020. Culture of marine shrimp (*Litopenaeus vannamei*) in biofloc technology system using artificially salinized freshwater: Zootechnical performance, economics and nutritional quality. *Aquaculture* 520, 734960. <https://doi.org/10.1016/j.aquaculture.2020.734960>
- Ponce-Palafox, J., Martinez-Palacios, C.A., Ross, L.G., 1997. The effects of salinity and temperature on the growth and survival rates of juvenile white shrimp, *Penaeus vannamei*, Boone, 1931. *Aquaculture* 157, 107–115. [https://doi.org/10.1016/s0044-8486\(97\)00148-8](https://doi.org/10.1016/s0044-8486(97)00148-8)
- Ramírez-Rochín, J., Frías-Espericueta, M.G., Fierro-Sañudo, J.F., Alarcón-Silvas, S.G., Fregoso-López, M.G., Páez-Osuna, F., 2017. Acute toxicity of nitrite on white shrimp *Litopenaeus vannamei* (Boone) juveniles in low-salinity water. *Aquac. Res.* 48, 2337–2343. <https://doi.org/10.1111/are.13069>
- Romano, N., Dauda, A.B., Ikhsan, N., Karim, M., Kamarudin, M.S., 2018. Fermenting rice bran as a carbon source for biofloc technology improved the water quality, growth, feeding efficiencies, and biochemical composition of African catfish *Clarias gariepinus* juveniles. *Aquac. Res.* 49, 3691–3701. <https://doi.org/10.1111/are.13837>
- Roy, L.A., Davis, A., 2010. Requirements for the culture of the Pacific white shrimp, *Litopenaeus vannamei*, reared in low salinity waters: water modification and nutritional strategies for improving production. *Avances En Nutrición Acuicola*. 61–78.

- Roy, L.A., Davis, D.A., Saoud, I.P., Henry, R.P., 2007. Effects of varying levels of aqueous potassium and magnesium on survival, growth, and respiration of the Pacific white shrimp, *Litopenaeus vannamei*, reared in low salinity waters. *Aquaculture* 262, 461–469. <https://doi.org/10.1016/j.aquaculture.2006.10.011>
- Roy, L. A., Davis, D. A., Saoud, I. P., Boyd, C. A., Pine, H. J., Boyd, C. E., 2010. Shrimp culture in inland low salinity waters. *Reviews in Aquaculture*, 2(4), 191–208. <https://doi.org/10.1111/j.1753-5131.2010.01036.x>
- Samocha, T.M., 2019. Sustainable Biofloc Systems for Marine Shrimp. Academic Press, San Diego, CA.
- Santos, R.B., Coelho Filho, P.A., Gonçalves, A.P., Santos, R.A., Lins Rodrigues, M., Souza Correia, E., Oliveira, V.Q., Brito, L.O., 2022. Effects of organic carbon sources on water quality, microbial flocs protein and performance of *Macrobrachium rosenbergii* post-larvae reared in biofloc and synbiotic systems. *Aquac. Res.* 53, 388–397. <https://doi.org/10.1111/are.15580>
- Saoud, I.P., Davis, D.A., Rouse, D.B., 2003. Suitability studies of inland well waters for *Litopenaeus vannamei* culture. *Aquaculture* 217, 373–383. [https://doi.org/10.1016/s0044-8486\(02\)00418-0](https://doi.org/10.1016/s0044-8486(02)00418-0)
- Schmidt-Nielsen, K., 1990. Animal physiology: adaption and environment, 4th ed. Cambridge University Press, Cambridge, New York.
- Van Wyk, P., Davis-Hodgkins, M., Laramore, R.W., Main, K.L., Mountain, J., Scarpa, J., 1999. Farming Marine Shrimp in Recirculating Freshwater Systems. Harbor Branch Oceanographic Institution, v 7, pp 125-140.

Zacarias, S., Schweitzer, R., Arantes, R., Galasso, H., Pinheiro, I., Espirito Santo, C.,

Vinatea, L., 2018. Effect of different concentrations of potassium and magnesium on performance of *Litopenaeus vannamei* postlarvae reared in low-salinity water and a biofloc system. J. Appl. Aquacult. 31, 1–12.

<https://doi.org/10.1080/10454438.2018.1536009>

Zhang, J., Liu, Y., Tian, L., Yang, H., Liang, G., Xu, D., 2012. Effects of dietary mannan oligosaccharide on growth performance, gut morphology and stress tolerance of juvenile Pacific white shrimp, *Litopenaeus vannamei*. Fish Shellfish Immunol. 33,

1027–1032. <https://doi.org/10.1016/j.fsi.2012.05.001>

Table 1. Percentage % increment of the target ion (after 72 h of the application of 100 g/m³) by chemical compounds used for ionic adjustment in intensive nursery synbiotic system.

Commercial product		MgCl ₂ . 6H ₂ O	CaCO ₃	KCl
%	pH ^a	0.17 ± 0.01	0.32 ± 0.06	0.41 ± 0.05
increment	Ca ²⁺	-	30.4 ± 0.73	-
of the	K ⁺	-	-	51.1 ± 11.38
target ion	Mg ²⁺	10.52 ± 1.37	-	-

Note: The data correspond to the mean of 2 replicates ± standard deviation. KCl: potassium chloride; CaCO₃: calcium carbonate; MgCl₂. 6 H₂O: magnesium chloride hexahydrate. ^apH in absolute value of the hydrogenic potential; Ca²⁺: calcium ion; K⁺: potassium ion; Mg²⁺: magnesium ion.

Table 2. Total ammonia nitrogen (TAN) concentration and non-ionized ammonia (NH₃-N) produced in the experimental units from NH₄Cl solution application.

	TAN (mg L ⁻¹)		N-NH ₃ (mg L ⁻¹)	
	SW	SWD, 1IA, 2IA, 3IA	SW	SWD, 1IA, 2IA, 3IA
Initial	2.40	4.65	0.31	0.31
24hrs	2.20	4.62	0.30	0.26
48hrs	2.50	4.60	0.39	0.37
72hrs	3.23	6.18	0.39	0.41

Note: Data correspond to the mean concentration of TAN and NH₃-N. SW (seawater - salinity 31); SWD (seawater diluted to a salinity of 2.3); 1IA (salinity 2.3- ionic adjustment to achieve Ca:Mg:K ratio of 1:3:1 only on the 1st day); 2IA (salinity 2.3 - ionic adjustment to achieve Ca:Mg:K ratio of 1:3:1 on the 1st and 20th day) and 3IA- (salinity 2.3- ionic adjustment to achieve Ca:Mg:K ratio of 1:3:1 on the 1st, 10th and 20th day).

Table 3. Zootechnical performance of *Litopenaeus vannamei* after 40 days in a synbiotic nursery system with different ionic adjustment frequencies in oligohaline water.

Variables	Treatments									
	SW		SWD		1IA		2IA		3IA	
Final weight (g)	1.02 ^B	± 0.06	1.16 ^A	± 0.04	1.18 ^A	± 0.02	1.15 ^A	± 0.05	1.14 ^A	± 0.10
Yield (Kg m⁻³)	1.73 ^B	± 0.08	2.13 ^A	± 0.05	2.10 ^A	± 0.20	2.13 ^A	± 0.15	2.15 ^A	± 0.15
Survival (%)	84.72 ^B	± 0.03	91.67 ^{AB}	± 1.67	88.61 ^{AB}	± 7.19	92.22 ^{AB}	± 2.92	94.44 ^A	± 2.68
FCR	1.02	± 0.05	0.97	± 0.02	0.99	± 0.10	0.97	± 0.07	0.96	± 0.06
SGR (% day⁻¹)	11.49 ^B	± 0.15	11.81 ^A	± 0.09	11.86 ^A	± 0.05	11.80 ^A	± 0.10	11.76 ^A	± 0.22

Data correspond to the mean (n=3) ± standard deviation. Different letters correspond to One-Way ANOVA, followed by Duncan's test ($p < 0.05$) among treatments. SW (seawater - salinity 31); SWD (seawater diluted to a salinity of 2.3); 1IA (salinity 2.3- ionic adjustment to achieve Ca:Mg:K ratio of 1:3:1 only on the 1st day); 2IA (salinity 2.3 - ionic adjustment to achieve Ca:Mg:K ratio of 1:3:1 on the 1st and 20th day) and 3IA- (salinity 2.3- ionic adjustment to achieve Ca:Mg:K ratio of 1:3:1 on the 1st, 10th and 20th day). FCR: feed conversion ratio; SGR: specific growth rate.

Table 4. Water quality variables measured during *Litopenaeus vannamei* synbiotic nursery system with different ionic adjustment frequencies in oligohaline water.

Variables	Treatments															
		SW			SWD			1IA			2IA			3IA		
DO** (mg L ⁻¹)	Mean ± SD	4.75 ^B	±	0.50	5.72 ^A	±	0.67	5.72 ^A	±	0.64	5.71 ^A	±	0.64	5.71 ^A	±	0.70
	Range	4.26	-	5.24	5.07	-	6.47	5.10	-	6.74	5.07	-	6.73	5.07	-	6.44
Temperature* (°C)	Mean ± SD	30.85	±	0.57	30.75	±	0.68	30.69	±	0.53	30.90	±	0.73	30.98	±	0.84
	Range	30.32	-	31.65	30.36	-	31.07	30.30	-	31.46	30.38	-	31.40	30.24	-	31.49
SS** (mL L ⁻¹)	Mean ± SD	0.08 ^B	±	0.15	0.63 ^{Bb}	±	1.14	3.95 ^{Aa}	±	4.42	3.96 ^{Aa}	±	3.97	2.23 ^{ABab}	±	2.90
	Range	0.00	-	0.28	0.00	-	2.00	0.70	-	13.67	0.10	-	12.67	0.30	-	11.00
Salinity* (g L ⁻¹)	Mean ± SD	31.17 ^A	±	1.05	2.26 ^{Bb}	±	0.06	2.39 ^{Bb}	±	0.06	2.59 ^{Ba}	±	0.17	2.74 ^{Ba}	±	0.19
	Range	29.98	-	32.82	2.15	-	2.33	2.32	-	2.48	2.37	-	2.93	2.47	-	2.93
pH*	Mean ± SD	8.13 ^A	±	0.20	8.04 ^{ABab}	±	0.22	7.96 ^{Bab}	±	0.29	7.90 ^{Bb}	±	0.33	7.98 ^{Bab}	±	0.28
	Range	7.98	-	8.35	7.82	-	8.36	7.59	-	8.31	7.37	-	8.38	7.48	-	8.28
TA** (mg CaCO ₃ L ⁻¹)	Mean ± SD	171.33 ^A	±	30.96	92.33 ^{Ba}	±	16.35	87.33 ^{BCab}	±	19.35	83.33 ^{Cab}	±	17.49	87.33 ^{BCb}	±	15.91

TH* (mg L ⁻¹)	Range	125.00	225.00	65.00	- 120.00	55.00	- 115.00	55.00	- 110.00	55.00	- 115.00
	Mean ± SD	6,160.00 ^A	± 104.31	490.13 ^{Ed}	± 33.19	586.67 ^{Dc}	± 61.59	710.13 ^{Cb}	± 216.32	801.07 ^{Ba}	± 238.88
PO ₄ ³⁻ * (mg L ⁻¹)	Range	5,960.00	- 6,240.00	452.00	- 588.00	452.00	- 648.00	452.00	- 988.00	452.00	- 1,088.00
	Mean ± SD	32.65 ^C	± 8.21	33.81 ^{BCb}	± 7.71	38.00 ^{Aa}	± 10.82	36.05 ^{ABab}	± 9.47	37.69 ^{Aa}	± 10.25
TAN* (mg L ⁻¹)	Range	23.83	- 45.36	23.00	- 47.08	23.00	- 52.05	23.00	- 49.40	23.00	- 50.82
	Mean ± SD	0.44 ^B	± 0.51	0.61 ^A	± 0.56	0.55 ^{AB}	± 0.50	0.55 ^{AB}	± 0.46	0.66 ^A	± 0.53
NO ₂ ⁻ -N** (mg L ⁻¹)	Range	0.00	- 1.45	0.04	- 1.81	0.00	- 1.54	0.07	- 1.61	0.24	- 1.66
	Mean ± SD	11.60 ^A	± 12.26	0.52 ^B	± 0.45	0.46 ^B	± 0.15	0.49 ^B	± 0.28	0.50 ^B	± 0.30
NO ₃ ⁻ -N* (mg L ⁻¹)	Range	1.20	- 33.00	0.22	- 1.60	0.32	- 0.76	0.20	- 1.30	0.20	- 1.40
	Mean ± SD	1.37	± 1.04	1.73	± 1.26	1.58	± 0.84	1.91	± 1.36	1.61	± 0.90
	Range	0.20	- 3.32	0.13	- 3.80	0.13	- 3.36	0.13	- 4.70	0.13	- 2.98

Data correspond to the mean (n=3) ± standard deviation. Different capital letters represent significant differences ($p < 0.05$) among the treatments (considering all treatments) and lowercase letters represent significant differences ($p < 0.05$) among low salinity treatments (excluding SW treatment). * Repeated measures ANOVA results followed by Duncan's test. ** Friedman test results followed by Dunn's test. SW (seawater - salinity 31); SWD (seawater diluted to a salinity of 2.3); 1IA (salinity 2.3- ionic adjustment to achieve Ca:Mg:K ratio of 1:3:1 only on the 1st day);

2IA (salinity 2.3 - ionic adjustment to achieve Ca:Mg:K ratio of 1:3:1 on the 1st and 20th day) and 3IA- (salinity 2.3- ionic adjustment to achieve Ca:Mg:K ratio of 1:3:1 on the 1st, 10th and 20th day). SD: standard deviation; DO: dissolved oxygen; SS: settleable solids; pH: hydrogen potential; TA: total alkalinity; TH: total hardness; PO_4^{3-} : orthophosphate; TAN: total ammonia nitrogen; NO_2^- -N: nitrite nitrogen; NO_3^- -N: nitrate nitrogen.

Table 5. Ion concentration (mg L⁻¹) in *Litopenaeus vannamei* culture water using different ionic adjustment frequencies in a synbiotic nursery system.

Variables	Treatments									
	SW		SWD		1IA		2IA		3IA	
Ca²⁺	402.7 ^A	± 39.7	38.0 ^B	± 6.7	37.5 ^B	± 3.9	46.9 ^B	± 12.2	51.3 ^B	± 11.2
Mg²⁺	1,252.3 ^A	± 21.7	96.0 ^E	± 6.2	119.8 ^D	± 15.7	144.1 ^C	± 46.4	163.5 ^B	± 52.0
K⁺	217.1 ^A	± 61.3	48.3 ^B	± 10.5	52.8 ^B	± 11.5	57.7 ^B	± 14.6	61.5 ^B	± 17.0
SO₄²⁻	2,817.5 ^A	± 339.0	231.9 ^B	± 23.6	242.9 ^B	± 30.9	258.9 ^B	± 52.7	294.4 ^B	± 65.2
Cl⁻	19,970.2 ^A	± 2,137.8	1,201.8 ^B	± 66.0	1,260.8 ^B	± 64.6	1,323.5 ^B	± 111.2	1,425.1 ^B	± 139.5
Na⁺	11,035.5 ^A	± 1,181.4	664.1 ^B	± 36.5	696.7 ^B	± 35.7	731.3 ^B	± 61.5	787.5 ^B	± 77.1
Mg:Ca	3.1:1 ^A		2.5:1 ^B		3.2:1 ^A		3.1:1 ^A		3.2:1 ^A	
Na:K	51.7:1 ^A		13.7:1 ^B		13.2:1 ^B		12.7:1 ^B		12.8:1 ^B	
Ca:Mg:K	1:3.1:0.5		1:2.5:1.3		1:3.2:1.4		1:3.1:1.2		1:3.2:1.2	

Data correspond to the mean (n=15) ± standard deviation. Different letters represent repeated measures ANOVA, followed by Duncan's test ($p < 0.05$). SW (seawater - salinity 31); SWD (seawater diluted to a salinity of 2.3); 1IA (salinity 2.3- ionic adjustment to achieve Ca:Mg:K ratio of

1:3:1 only on the 1st day); 2IA (salinity 2.3 - ionic adjustment to achieve Ca:Mg:K ratio of 1:3:1 on the 1st and 20th day) and 3IA- (salinity 2.3- ionic adjustment to achieve Ca:Mg:K ratio of 1:3:1 on the 1st, 10th and 20th day). Ca^{2+} : calcium ion; Mg^{2+} : magnesium ion; K^{+} : potassium ion; SO_4^{2-} : sulfate ion; Cl^{-} : chloride ion; Na^{+} : sodium ion; Mg:Ca: ratio of magnesium and calcium ions; Na:K: ratio of sodium and potassium ions; Ca:Mg:K: ratio of calcium, magnesium and potassium ions.

Table 6. Microbial floc mineral concentration (g Kg⁻¹; dry weight) in the *Litopenaeus vannamei* synbiotic nursery using different ionic adjustment frequencies in oligohaline water.

Minerals (microbial floc)	Treatments														
	SW			SWD			1IA			2IA			3IA		
P (g Kg⁻¹)	4.25	±	1.07	2.34	±	0.47	1.90	±	0.10	2.05	±	0.20	1.87	±	0.29
K (g Kg⁻¹)	0.63 ^A	±	0.10	0.15 ^B	±	0.01	0.19 ^B	±	0.02	0.22 ^B	±	0.01	0.24 ^B	±	0.03
Ca (g Kg⁻¹)	20.15 ^A	±	3.84	8.88 ^B	±	1.79	10.02 ^{AB}	±	1.21	11.78 ^{AB}	±	0.59	9.40 ^{AB}	±	1.32
Mg (g Kg⁻¹)	1.52 ^A	±	0.10	0.38 ^D	±	0.06	0.44 ^{CD}	±	0.03	0.52 ^{BC}	±	0.03	0.58 ^B	±	0.03
Na (g Kg⁻¹)	5.73 ^A	±	1.13	0.63 ^B	±	0.06	0.61 ^B	±	0.08	0.69 ^B	±	0.04	0.68 ^B	±	0.01
S (g Kg⁻¹)	0.94 ^A	±	0.08	0.32 ^B	±	0.07	0.31 ^B	±	0.03	0.36 ^B	±	0.07	0.40 ^B	±	0.02
Cu (mg Kg⁻¹)	7.05 ^A	±	0.35	3.87 ^B	±	1.08	3.63 ^B	±	0.71	3.99 ^B	±	0.35	4.27 ^B	±	0.18
Fe (mg Kg⁻¹)	1,167.46	±	50.66	827.19	±	253.28	621.65	±	54.86	703.45	±	79.40	672.55	±	103.28
Mn (mg Kg⁻¹)	11.31	±	1.52	11.85	±	2.42	13.94	±	4.77	16.29	±	4.68	15.02	±	2.30
Zn (mg Kg⁻¹)	73.70 ^A	±	10.66	41.02 ^B	±	7.70	38.50 ^B	±	3.34	37.06 ^B	±	12.38	39.67 ^B	±	5.67

Data correspond to the mean ($n = 3$) $\pm$ standard deviation. Different letters represent One-Way ANOVA, followed by Duncan's test ($p < 0.05$). SW (seawater - salinity 31); SWD (seawater diluted to a salinity of 2.3); 1IA (salinity 2.3- ionic adjustment to achieve Ca:Mg:K ratio of 1:3:1 only on the 1st day); 2IA (salinity 2.3 - ionic adjustment to achieve Ca:Mg:K ratio of 1:3:1 on the 1st and 20th day) and 3IA- (salinity 2.3- ionic adjustment to achieve Ca:Mg:K ratio of 1:3:1 on the 1st, 10th and 20th day). P: phosphor; K: potassium; Ca: calcium; Mg: magnesium; Na: sodium; S: sulfur; Cu: copper; Fe: iron; Mn: manganese; Zn: zinc.

Table 7. Marine shrimp *Litopenaeus vannamei* biomass mineral concentration (g Kg⁻¹; dry weight) using different ionic adjustment frequencies in a synbiotic nursery system.

Minerals	Treatments														
(Shrimp)	SW			SWD			1IA			2IA			3IA		
P (g Kg ⁻¹)	12.54	±	0.06	13.31	±	0.58	12.31	±	0.56	12.67	±	0.32	12.60	±	1.27
K (g Kg ⁻¹)	12.03	±	0.11	13.07	±	0.73	13.10	±	1.00	12.35	±	0.76	11.44	±	1.21
Ca (g Kg ⁻¹)	45.88	±	4.19	42.92	±	4.12	55.61	±	6.03	48.83	±	0.85	51.80	±	12.31
Mg (g Kg ⁻¹)	3.90 ^A	±	0.04	2.87 ^B	±	0.17	2.87 ^B	±	0.03	3.13 ^{AB}	±	0.02	3.26 ^{AB}	±	0.12
Na (g Kg ⁻¹)	11.91 ^A	±	0.47	10.40 ^{AB}	±	0.70	9.79 ^{AB}	±	0.49	9.38 ^B	±	0.34	9.67 ^{AB}	±	0.01
S (g Kg ⁻¹)	6.27 ^{AB}	±	0.84	6.31 ^{AB}	±	0.72	5.11 ^B	±	0.54	6.95 ^{AB}	±	0.03	7.69 ^A	±	0.76
Cu (mg Kg ⁻¹)	9.83	±	1.39	11.98	±	0.60	15.10	±	0.83	15.75	±	2.81	15.28	±	1.14
Fe (mg Kg ⁻¹)	102.52 ^B	±	3.33	226.85 ^A	±	14.68	226.50 ^A	±	37.44	235.38 ^A	±	40.20	182.80 ^A	±	53.03
Mn (mg Kg ⁻¹)	3.95 ^B	±	0.33	7.20 ^A	±	0.44	7.05 ^A	±	1.52	6.88 ^A	±	1.52	7.85 ^A	±	0.23
Zn (mg Kg ⁻¹)	61.22 ^B	±	1.38	64.50 ^{AB}	±	1.18	64.30 ^{AB}	±	0.85	66.77 ^A	±	3.03	62.90 ^{AB}	±	3.40

Data correspond to the mean (n=3) ± standard deviation. Different letters represent One-Way ANOVA, followed by Duncan's test (p < 0.05). SW (seawater - salinity 31); SWD (seawater diluted to a salinity of 2.3); 1IA (salinity 2.3- ionic adjustment to achieve Ca:Mg:K ratio of 1:3:1 only on the 1st day); 2IA (salinity 2.3 - ionic adjustment to achieve Ca:Mg:K ratio of 1:3:1 on the 1st and 20th day) and 3IA- (salinity 2.3- ionic adjustment

to achieve Ca:Mg:K ratio of 1:3:1 on the 1st, 10th and 20th day). P: phosphor; K: potassium; Ca: calcium; Mg: magnesium; Na: sodium; S: sulfur; Cu: copper; Fe: iron; Mn: manganese; Zn: zinc.

3. REFERENCIA BIBLIOGRÁFICA

- APHA, 2005. Standard Methods for the Examination of Water and Wastewater, 21st Ed. American Public Health Association/American Water Works Association/Water Environment Federation, Washington, USA.
- APHA, 2012. Standard Methods for the Examination of Water and Wastewater, 22th Ed. American Public Health Association, Washington, USA.
- Avnimelech, Y., 2012. Biofloc Technology - A Practical Guide Book, 2nd Ed. The World Aquaculture Society, Baton Rouge, Louisiana, United States.
- Boyd, C. E., 2002. Anion-cation balance in water. Global Aquaculture Advocate 74-75. <https://www.aquaculturealliance.org/advocate/anion-cation-balance-in-water/?headlessPrint=AAAAPIA9c8r7gs82oWZBA>.
- Boyd, C.E, 2020. Water quality: an introduction. 3th Ed. Springer Science, Switzerland.
- Boyd, C.E., Tucker, C. S., 1998. Pond aquaculture water quality management/ Claude E. Boyd, Craig S. Tucker. Kluwer Academic Publishers, Massachusetts, EUA.
- Boyd, C.E., Thunjai, T., 2003. Concentrations of major ions in waters of inland shrimp farms in China, Ecuador, Thailand, and the United States. Journal of the World Aquaculture Society, 34, 524-532.
- Boyd, Claude E., and Craig S. Tucker. Pond aquaculture water quality management. Springer Science & Business Media, 2012.
- Boyd, C., Somridhivej, B., Tucker, C.S., 2016. Alkalinity and Hardness: Critical but Elusive Concepts in Aquaculture. Journal of the World Aquaculture Society, 47(1):6-41.
- Boyd, Claude E. Water quality: an introduction. Springer Nature, 2019.
- Brito, L.O., Santos, I.G.S., Abreu, J.L., Araújo, M.T., Severi, W., Gálvez, A.O., 2016. Effect of the addition of diatoms (*Navicula spp.*) and rotifers (*Brachionus plicatilis*) on water quality and growth of the *Litopenaeus vannamei* postlarvae reared in a biofloc system. Aquaculture Research, 47, 3990–3997.
- Burbano-Gallardo, E., Imues-Figueroa, M.A., Gonzalez-Legarda, E.A., Brito, L.O., Galvez, A.O., Vinatea Arana, L.A., 2015. Survival of *Litopenaeus vannamei* postlarvae under osmotic stress and its relation with the molt stage. Revista de Biología Marina y Oceanografía, Chile, 50, 2, 323-330.

- Casillas-Hernández, R., Magallón-Barajas, F., Portillo-Clarck, G., Páez-Osuna, F., 2006. Nutrient mass balances in semi-intensive shrimp ponds from Sonora, Mexico using two feeding strategies: Trays and mechanical dispersal. *Aquaculture*, 258(1-4), 289–298.
- Carmouze, J., 1994. O metabolismo dos ecossistemas aquáticos – Fundamentos teóricos, métodos de estudo e análises químicas. São Paulo: Edgard Blücher. 254 p.
- Cheng, K.-M., Hu, C.-Q., Liu, Y.-N., Zheng, S.-X, Qi, X.-J, 2005. Dietary magnesium requirement and physiological responses of marine shrimp *Litopenaeus vannamei* reared in low salinity water. *Aquaculture Nutrition*, 11 (5), 385-393.
- Correia, E.S., Wilkenfeld, J.S., Morris, T.C., Wei, L., Prangnell, D.I., Samocha, T.M., 2014. Intensive nursery production of the pacific white shrimp *Litopenaeus vannamei* using two commercial feeds with high and low protein content in a biofloc-dominated system. *Aquacultural engineering*. 59, 48 – 54.
- Custodio, E., Llamas, M.R., 1983. Hidrología subterránea, 2nd ed. Ediciones Omega, Barcelona.
- Da Silva, K.R., Wasielesky Jr., W., Abreu, P.C., 2013. Nitrogen and phosphorus in the biofloc production of the pacific white shrimp, *Litopenaeus vannamei*. *Journal of the World Aquaculture Society* 44, 1.
- Davis, D.A., Lawrence, A.L., 1997. Minerals. In: D'Abramo, L.R., Conklin, D.E., Akiyama, D.M., eds. *Crustacean Nutrition*, 6, 150–163.
- Ebeling, J.M., Timmons, M.B., Bisogni, J.J., 2006. Engineering analysis of the stoichiometry of photoautotrophic, autotrophic, and heterotrophic removal of ammonia–nitrogen in aquaculture systems. *Aquaculture*, 257 (1-4), 346–358.
- Egna, H.S., Boyd, C.E., 1997. Dynamics of pond aquaculture. Boca Raton, Florida, USA: CRC Press LLC.
- Eldyasti, A., Nakhla, G., Zhu, J., 2013. Impact of calcium on biofilm morphology, structure, detachment and performance in denitrifying fluidized bed bioreactors (DFBBRs). *Chemical Engineering Journal* 232, 183–195.
- Elser, J.J., Dobberfuhl, D.R., MacKay, N.A., Schampel, J.H., 1996. Organism size, life history, and N:P stoichiometry. *Bioscience* 46, 674–684.
- Esparza-Leal, Héctor Manuel, et al. "The effect of low salinity water with different ionic composition on the growth and survival of *Litopenaeus vannamei* (Boone, 1931) in intensive culture." *Journal of Applied Aquaculture* 21.4 (2009): 215-227.

- Esparza-Leal, H.M., Xavier, J. A.A., Wasielesky, W., 2016. Performance of *Litopenaeus vannamei* postlarvae reared in indoor nursery tanks under biofloc conditions at different salinities and zero-water exchange. *Aquaculture international*, 24, 5, 1435-1447.
- Esteves, F.A., 2011. Fundamentos de Limnologia. 3 rd. Rio de Janeiro: Interciência. 826p.
- Ferreira, G.S., Fernandes Silva, V., Aranha Martins, M., da Silva, A.C.C.P., Machado, C., Quadros Seiffert, W., Vieira, F.N., 2020. Strategies for ammonium and nitrite control in *Litopenaeus vannamei* nursery systems with bioflocs. *Aquacultural Engineering* 88, 102040.
- Fox, J.; Weisberg, S., 2019. An {R} Companion to Applied Regression, Third Edition. Thousand Oaks CA: Sage. URL: <https://socialsciences.mcmaster.ca/jfox/Books/Companion/>
- Fries, J., 1971. Análisis de Trazas. Métodos Fotometricos Comprobados. Darmstadt: Merck. 130p.
- Fries, J., Getrost, H., 1977. Organic Reagents for Trace Analysis. Darmstadt, Germany: E. Merck. 453p.
- Furtado, P.S., Poersch, L.H., Wasielesky, W., 2015. The effect of different alkalinity levels on *Litopenaeus vannamei* reared with biofloc technology (BFT). *Aquaculture International*, 23, 345-358.
- Godwin, C.M., Whitaker, E.A., Cotner, J.B., 2017. Growth rate and resource imbalance interactively control biomass stoichiometry and elemental quotas of aquatic bacteria. *Ecology*, 98(3), 820–829.
- Gosling, E., 2003. Marine Bivalve Molluscs: biology, ecology and culture. Wiley-Blackwell, UK. 456p.
- Gross, A.; Abutbul, S.; Zilberg, D., 2004. Acute and chronic effects of nitrite on white shrimp *Litopenaeus vannamei* cultured in low salinity brackish water. *J World Aquac Soc* 35:315–321.
- Harrell Jr, F.E., 2020. with contributions from Charles Dupont and many others. Hmisc: Harrell Miscellaneous. R package version 4.4-1. <https://CRAN.Rproject.org/package=Hmisc>
- Jung, J.Y., Damusaru, J.H., Park, Y., Kim, K., Seong, M., Je, H-W., Kim, S., Bai, S. C., 2017. Autotrophic biofloc technology system (ABFT) using *Chlorella vulgaris* and

- Scenedesmus obliquus* positively affects performance of Nile tilapia (*Oreochromis niloticus*). *Algal Research*, 27, 259–264.
- Khanjani, Mohammad Hossein, Morteza Alizadeh, and Moslem Sharifinia. "Rearing of the Pacific white shrimp, *Litopenaeus vannamei* in a biofloc system: The effects of different food sources and salinity levels." *Aquaculture Nutrition* 26.2 (2020): 328-337.
- Khanjani, Mohammad Hossein, Moslem Sharifinia, and Saeed Hajirezaee. "Effects of different salinity levels on water quality, growth performance and body composition of Pacific white shrimp (*Litopenaeus vannamei* Boone, 1931) cultured in a zero water exchange heterotrophic system." *Annals of Animal Science* 20.4 (2020): 1471-1486.
- Laramore, S., Laramore, C.R., Scarpa, J., 2001. Effect of Low Salinity on Growth and Survival of Postlarvae and Juvenile *Litopenaeus vannamei*. *Journal of the world aquaculture society*, 32, 4.
- Lin, Y.C., Chen, J.C., 2001. Acute toxicity of ammonia on *Litopenaeus vannamei* Boone juveniles at different salinity levels. *Journal of Experimental Marine Biology and Ecology*, 259(1), 109–119.
- Lin, Y.C., Chen, J.C. 2003. Acute toxicity of nitrite on *Litopenaeus vannamei* (Boone) juveniles at different salinity levels. *Aquaculture*, 224(1-4), 193–201. doi:10.1016/s0044-8486(03)00220-5
- Luo, G., Liang, W., Tan, H., Yao, C., Zhang, N., Lu, L., 2013. Effects of calcium and magnesium addition on the start-up of sequencing batch reactor using biofloc technology treating solid aquaculture waste. *Aquacultural Engineering*, 57, 32-37.
- Mackereth, J. F. H., Heron, J., Talling, J. F., 1978. Water analysis: some revised methods for limnologists. *Freshwater Biological Association*, 36, 121.
- Marinho, Y.F., Brito, L.O., Campos, C.V.F.S., Severi, W., Andrade, H.A., Galvez, A.O., 2016. Effect of the addition of *Chaetoceros calcitrans*, *Navicula* sp. and *Phaeodactylum tricornutum* (diatoms) on phytoplankton composition and growth of *Litopenaeus vannamei* (Boone) postlarvae reared in a biofloc system. *Aquaculture Research*, 48, 4155–4164.
- McNevin, A.A., Boyd, C.E., Silapajarn, O., Silapajarn, K., 2004. Ionic supplementation of pond waters for inland culture of marine shrimp. *Journal of World Aquaculture Society*. 35, 460-467.

- Naik, P., 2012. Essentials of Biochemistry. Nova Dheli: Jaypee Brothers Medical Publishers. 451 p.
- Nehru, E., Rao, A.C., Pamanna, D., Ranjith, P., Lokesh, B., 2018a. Effect of Aqueous Minerals Supplementation on Growth and Survival of *Litopenaeus vannamei* in Low Salinity Water. International Journal of Current Microbiology and Applied Sciences, 7, 1, 1706-1713.
- Nehru, E., Chandrasekhara Rao, A., Pamanna, D., priyadarshini, N., Anil Kumar, P., Raveendra, M., 2018b. Effect of Dietary Minerals Supplementation on Growth and Survival of *Litopenaeus vannamei* in Low Salinity Water. International Journal of Current Microbiology and Applied Sciences, 7, 3, 3040-3049.
- Pimentel, O.A.L.F., Amado, A.M., They, N.H., 2020. Higher nitrogen and phosphorus immobilization in bioflocs is associated with higher temperature and increased suspended solids in shrimp farming with biofloc technology. Aquaculture Research. 51, 3888–3899.
- Pimentel, O. A. L. F., Oliveira, V. Q., Oliveira, C. R. do R., Severi, W., Olivera Gálvez, A., Amado, A. M., & Brito, L. O. (2021). Assessment of different ionic adjustment strategies in low-salinity water on the growth of *Litopenaeus vannamei* and microbial community stoichiometry in a synbiotic nursery system. Aquaculture Research. doi:10.1111/are.15552
- Pohlert T., 2014. _The Pairwise Multiple Comparison of Mean Ranks Package (PMCMR)_. R package, <URL: <https://CRAN.R-project.org/package=PMCMR>>.
- R Core Team., 2020. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.
- Ramírez-Rochín, J., Frías-Espéricueta, M.G., Fierro-Sañudo, J.F., Alarcón-Silvas, S.G., Fregoso-López, M.G., Páez-Osuna, F., 2016. Acute toxicity of nitrite on white shrimp *Litopenaeus vannamei* (Boone) juveniles in low-salinity water. Aquaculture Research, 48(5), 2337–2343.
- Romano, N., Dauda, A.B., Ikhsan, N., Karin, M., Kamarudin, M.S., 2018. Fermenting rice bran as a carbon source for biofloc technology improved the water quality, growth, feeding efficiencies, and biochemical composition of African catfish *Clarias gariepinus* juveniles. Aquaculture research, 49, 3691-3701.

- Roy, L.A., Davis, D.A., Saoud, I.P., Henry, R.P., 2007. Effects of varying levels of aqueous potassium and magnesium on survival, growth, and respiration of the Pacific white shrimp, *Litopenaeus vannamei*, reared in low salinity waters. *Aquaculture* 262, 461– 69.
- Roy, L.A., Davis, D.A., Saoud, I.P., Boyd, C.A., Pine, H.J., Boyd, C.E., 2010. Shrimp culture in inland low salinity waters. *Reviews in Aquaculture*, 2, 4, 191-208.
- Roy, L. A., Davis, D. A.; Whitis, G. N., 2009. Pond-to-pond variability in post-larval shrimp, *Litopenaeus vannamei*, survival and growth in inland low salinity waters of west Alabama. *Aquaculture Research*, 40, 1823-1829.
- Roy, L.A., Davis, D.A., 2010. Requirements for the culture of the Pacific white shrimp *Litopenaeus vannamei*, reared in low salinity waters: water modification and nutritional strategies for improving production. In: Cruz-Suárez, L.E., Ricque-Marie, D., Tapia-Salazar, M., Nietolópez, M.G., Villarreal-Cavazos, D.A., Gamboa-Delgado, J. (eds). *Avances en Nutrición Acuícola X - Memorias del X Simposio Internacional de Nutrición Acuícola*. 61-78.
- Sá, M.V.C., 2012. *Limnicultura: limnologia para aquicultura*. Fortaleza: Editora da Universidade Federal do Ceará. 218p.
- Samocha, T.M., Davis, A.D., Lawrence, A.L., Collins, C.R., Van Wyk, P., 2001. Intensive and superintensive production of the Pacific white *Litopenaeus vannamei* in greenhouse-enclosed raceway systems. *Book of Abstracts, Aquaculture*. Lake Buena Vista, Florida.
- Samocha, T.M., 2019. *Sustainable Biofloc Systems for Marine Shrimp – The Texas A&M AgriLife Research Experience*. Baton Rouge: The World Aquaculture Society. 463p.
- Silva Neto, S.R., Lavander, H.D., Souza, A.B., Cardoso Junior, L.O., Brito, L.O., Gálvez, A.O., 2012. Utilização de conchas de *Anomalocardia brasiliiana* como substrato de biofiltros no tratamento dos efluentes de cultivo heterotrófico de camarão marinho em laboratório. *Pesquisa Agropecuária Pernambucana*, Recife, 17, 20-23.
- Sterner, R.W., Elser, J.J., 2002. *Ecological Stoichiometry: The Biology of Elements from Molecules to the Biosphere*. United States of America: Princeton University Press. 464p.

- Tacon, A.J., 1987. The nutrition and feeding of farmed fish and shrimp. 1. The essential nutrients. Training Manual. Food and Agriculture Organization. Brasilia, Brazil, 73-84.
- Wei, T., Simko, V.R., 2017. package "corrplot": Visualization of a Correlation Matrix (Version 0.84). Available in: <https://github.com/taiyun/corrplot>.
- Valencia-Castañeda, G., Frías-Espéricueta, M.G., Vanegas-Pérez, R.C., Pérez-Ramírez, J.A., Chávez-Sánchez, M.C., Páez-Osuna, F., 2018. Acute Toxicity of Ammonia, Nitrite and Nitrate to Shrimp *Litopenaeus vannamei* postlarvae in Low-Salinity Water. Bulletin of Environmental Contamination and Toxicology, 101(2), 229–234.
- Valenzuela-Madrigal, I.E., Valenzuela-Quiñónez, W., Leal, H.M.E., Quiroz, G.R., Noriega, E.A.A., 2017. Effects of ionic composition on growth and survival of white shrimp *Litopenaeus vannamei* culture at low-salinity well water. Revista de biología marina y oceanografía, 52,1, 103-112.
- Van Wyk, P., Davis-Hodgkins, M., Laramore, R., Main, K.L., Mounta, J., Scarpa, J. 1999. Farming Marine Shrimp in Recirculating Freshwater Systems. Florida Department of Agriculture and Consumer Services – Harbor Branch Oceanic Institute, Tallahassee, Flórida. 125–140.
- Wetzel, R.G., Likens, G.E., 2000. Limnological analyses, 3rd ed. New York, NY: Springer Science + Business Media Inc.
- Zacarias, S., Schweitzer, R., Arantes, R., Galasso, H., Pinheiro, I., Espirito Santo, C., Vinatea, L., 2018. Effect of different concentrations of potassium and magnesium on performance of *Litopenaeus vannamei* postlarvae reared in low-salinity water and a biofloc system. Journal of Applied Aquaculture, 1-12.
- Zhang, J., Liu, Y., Tian, L., Yang, H., Liang, G., Xu, D., 2012. Effects of dietary mannan oligosaccharide on growth performance, gut morphology and stress tolerance of juvenile Pacific white shrimp, *Litopenaeus vannamei*. Fish & Shellfish immunology, 33, 4,1027-1032.