VICTOR ANDRADE DA SILVA

CRESCIMENTO, RESPOSTA FISIOLÓGICA E QUALIDADE DE FILÉ DE JUVENIS DE BEIJUPIRÁ (*Rachycentron canadum*) SUBMETIDOS A DIFERENTES DENSIDADES DE ESTOCAGEM

RECIFE

2013

UNIVERSIDADE FEDERAL RURAL DE PERNAMBUCO

PRÓ-REITORIA DE PESQUISA E PÓS-GRADUAÇÃO

PROGRAMA DE PÓS-GRADUAÇÃO EM RECURSOS PESQUEIROS E AQUICULTURA

CRESCIMENTO, RESPOSTA FISIOLÓGICA E QUALIDADE DE FILÉ DE JUVENIS DE BEIJUPIRÁ (*Rachycentron canadum*) SUBMETIDOS A DIFERENTES DENSIDADES DE ESTOCAGEM

Victor Andrade da Silva

Dissertação apresentada ao Programa de Pós-Graduação em Recursos Pesqueiros e Aquicultura da Universidade Federal Rural de Pernambuco como exigência para obtenção do título de Mestre.

Prof. Dr. Ronaldo Olivera Cavalli
Orientador

RECIFE

Fevereiro de 2013

Ficha Catalográfica

S586c Silva, Victor Andrade da
Crescimento, resposta fisiológica e qualidade de filé de juvenis de beijupirá (*Rachycentron canadum*) submetidos a diferentes densidades de estocagem / Victor Andrade da Silva. -- Recife, 2013.
59 f.: il.

Orientador (a): Ronaldo Oliveira Cavalli.
Dissertação (Mestrado em Recursos Pesqueiros e Aquicultura) – Universidade Federal Rural de Pernambuco, Departamento de Pesca e Aquicultura, Recife, 2013.
Referência.

1. Beijupirá 2. Peixe marinho 3. Piscicultura 4.Estresse crônico 5. Densidade de estocagem 6. Hematologia 7.Crescimento 8. Qualidade de filé I. Cavalli, Ronaldo Oliveira, Orientador II. Título

CDD 369.3

UNIVERSIDADE FEDERAL RURAL DE PERNAMBUCO

PRÓ-REITORIA DE PESQUISA E PÓS-GRADUAÇÃO

PROGRAMA DE PÓS-GRADUAÇÃO EM RECURSOS PESQUEIROS E AQUICULTURA

CRESCIMENTO, RESPOSTA FISIOLÓGICA E QUALIDADE DE FILÉ. DE JUVENIS DE BEIJUPIRÁ (*Rachycentron canadum*) SUBMETIDOS A DIFERENTES DENSIDADES DE ESTOCAGEM

Victor Andrade da Silva

Dissertação julgada adequada para obtenção do título de mestre em Recursos Pesqueiros e Aquicultura. Defendida e aprovada em 27/02/2013 pela seguinte Banca Examinadora.

Prof. Dr. Ronaldo Olivera Cavalli
(Orientador)
Departamento de Pesca e Aquicultura
Universidade Federal Rural de Pernambuco

Prof. Dr. Alfredo Olivera Gálvez
Departamento de Pesca e Aquicultura
Universidade Federal Rural de Pernambuco

Prof. Dr. Silvio Ricardo M. Peixoto
Departamento de Pesca e Aquicultura
Universidade Federal Rural de Pernambuco

Prof. Dr. Vinícius Ronzani Cerqueira
Departamento de Aquicultura
Universidade Federal de Santa Catarina

Profa. Dra. Roberta Borda Soares
Departamento de Pesca e Aquicultura
Universidade Federal Rural de Pernambuco

Dedicatória

A Deus, aos meus pais Jerônimo Gomes e Amara Andrade e a todos os meus familiares por terem me conduzido a um caminho de grandes realizações e de felicidades

Agradecimentos

Agradeço primeiramente a Deus, por permanecer comigo durante todos os momentos felizes e difíceis da minha vida e por guiar os meus passos durante a realização deste projeto.

Aos meus pais Jerônimo Gomes da Silva e Amara Andrade da Silva por estarem sempre presentes, pela amizade, carinho, amor e educação, que me foram oferecidos desde os primeiros dias de minha vida. A meu irmão, Kleber Andrade da Silva, que sempre tive confiança e amizade, pelo apoio durante mais uma etapa da minha vida. E a todos os familiares que participaram direta e indiretamente de minha formação moral e acadêmica.

À Universidade Federal Rural de Pernambuco e ao Programa de Pós-Graduação em Recursos Pesqueiros e Aquicultura pela oportunidade de obtenção do título de Mestre. E A todos os professores pelo aprendizado e trocas de experiência durante o tempo de formação.

Em especial ao professor e orientador Dr. Ronaldo O. Cavalli que desde a época de minha graduação contribui para minha formação profissional por meio de ensino e orientação qualificada, os quais persistiram em todos os momentos da realização desse projeto. E principalmente, por todo incentivo e oportunidades.

Aos membros da banca Dr. Alfredo Olivera Gálvez, Dr. Silvio Ricardo M. Peixoto, Dr. Vinícius Ronzani Cerqueira e a Drª. Roberta Borda Soares, pelas críticas e contribuições que enriqueceram todas as informações contidas nesta dissertação.

A todos do Laboratório de Piscicultura Marinha – LPM, em especial a Carolina, Cecília, Clebson, Ernesto, Gabriel, Larissa, Leilane, Reginaldo, Ricardo, Roberta, Santiago e Thales pelos momentos de descontração e amizade, e principalmente pelas trocas de conhecimento científico.

Aos pesquisadores Dr. Michael Schwarz e Drª. Jesse Trushenski pelo apoio e trabalho conjunto para a realização deste projeto. Aproveito a oportunidade para agradecer a todos os membros e professores do *Virginia Seafood Agricultural Research and Extension Center*, um dos laboratórios de pesquisa da Universidade *Virginia Tech*, onde foi realizado os ensaios experimentais dessa dissertação e principalmente ao gerente Steve Urick e a todos colegas de estagio pela amizade e apoio.

A Fundação de Amparo à Ciência e Tecnologia de Pernambuco - FACEPE pelo provimento de bolsa de pós-graduação e passagens aéreas para os Estados unidos nas modalidades IBPG e AMD, respectivamente. Nas quais contribuíram decisivamente para a realização dessa dissertação.

E, finalmente, a todos que contribuíram de alguma forma direta ou indireta na minha formação acadêmica e na realização desta dissertação.

Resumo

A densidade de estocagem é amplamente reconhecida como um fator crítico em aquicultura, uma vez que altas densidades representam uma potencial fonte de estresse crônico. Este estudo avaliou a resposta fisiológica, desempenho zootécnico e qualidade de filé de juvenis de beijupirá (*Rachycentron canadum*) submetidos a um estressor crônico, como altas densidades, visando à definição de condições de criação que levem à melhoria do bem estar animal e ao incremento da produtividade e qualidade do produto final. Os peixes foram testados por um período de 42 dias, em que juvenis de 46,32 ± 0,22 g (média ± erro padrão) foram aleatoriamente distribuídos em três densidades (3,30 ± 0,02; 6,67 ± 0,03; 13,15 ± 0,16 g L^{-1}), e em três repetições. Os tratamentos foram designados como baixa densidade (LD; 5 peixe/tanque), densidade média (MD; 10 peixe/tanque), e alta densidade (HD; 20 peixes/tanque), respectivamente. As concentrações de cortisol e lactato e a osmolalidade do plasma sanguíneo não foram afetadas pela densidade de estocagem, e seus valores estavam de acordo com as faixas basais previamente relatados para juvenis de beijupirá. No entanto, o ganho de peso e a taxa de crescimento específico foram significativamente maiores no grupo LD. A densidade de estocagem também influenciou o início e o desenvolvimento do rigor mortis. Juvenis de beijupirá criados na alta densidade apresentaram menor taxa de crescimento e qualidade de filé. Os resultados sugerem que juvenis de beijupirá criados nas altas densidades foram capazes de contrapor o estresse ao nível de composição do plasma. Contudo, energia da dieta e/ou reservas do corpo podem ter sido desviadas do crescimento como resposta à alta demanda metabólica causada pelo estresse. Dessa forma, o presente estudo sugere que juvenis de beijupirá devem ser mantidos em uma biomassa final de até 15 kg m^{-3}, aproximadamente para garantir máximo crescimento e melhor qualidade do produto final. Aparentemente, este estudo é o primeiro a abordar a influencia de um estresse crônico na resposta fisiológica e qualidade de filé de juvenis de beijupirá. Os resultados podem fornecer informações para avanços na criação dessa espécie, conduzindo a melhorias no bem estar animal, na qualidade do produto final e no aumento da produtividade.

Palavras-chave: Estresse crônico, rigor mortis, exsudação, qualidade de filé.

Abstract

Stocking density is recognized as a critical factor in aquaculture, since high densities represent a potential source of chronic stress. This study investigated the effects of a chronic stressor, such as high stocking densities, on growth, physiological responses, and flesh quality of juvenile cobia (*Rachycentron canadum*) tested for a period of 42 days. The trial was conducted in a recirculating aquaculture system equipped with tanks of 70 L tanks. Cobia of 46.32 ± 0.22 g (mean ± SEM) were randomly distributed into three stocking densities (3.30 ± 0.02; 6.67 ± 0.03; 13.15 ± 0.16 g L-1), and three replicates. Treatment groups were nominally assigned as low density (LD; 5 fish/tank), medium density (MD; 10 fish/tank), and high density (HD; 20 fish/tank), respectively. Plasma concentrations of cortisol, osmolality and lactate were not affected by density levels, and values were according with those reported as basal for juvenile cobia. However, growth rate were influenced by density levels, in which weight gain and specific growth rate were significantly higher in the LD group. Drip loss did not differ among density groups. However, stocking density influenced the onset and development of rigor mortis. Our findings indicated that rearing juvenile cobia at the high densities negatively affected growth rate and flesh quality. Results suggest that juvenile cobia reared at the high densities were able to counteract stress. However, energy from the diet and/or body reserves was diverted from growth as a response to the elevated metabolic demand caused by stress. In this regard, juvenile cobia should be maintained at a final biomass up to approximately 15 kg m^{-3} to ensure a maximal growth rate and an improved final product quality. To our knowledge, this is the first report assessing the influence of a chronic stressor on both physiological responses and flesh quality parameters in cobia. This may provide information to the advancement of cobia aquaculture, leading to improvements in fish welfare, final product quality and increased productivity.

Keywords: Chronic stress, rigor mortis, drip loss, flesh quality

Lista de figuras

Página

Figure 1- Mean (± SEM) drip loss (%) from flesh samples obtained from cobia (*Rachycentron canadum*) reared at low (LD), medium (MD) and high stocking densities (HD). There were not statistical differences (p = 0.057) in drip loss among density groups based on one-way ANOVA……..…… 47

Figure 2. Mean (± SEM) rigor indices (%) for juvenile cobia ((*Rachycentron canadum*) raised at low (LD), medium (MD) and high stocking densities (HD). Means with different letters are significantly different within each time point based on one-way ANOVA ($p<0.05$) with post hoc Duncan's multiple-range test……..……. 48

Lista de tabelas

Página

Table 1. Mean (± SEM) production performance parameters for cobia (*Rachycentron canadum*) reared at low (LD), medium (MD) and high densities (HD). Means with different letters are significantly different based on one-way ANOVA ($p<0.05$) followed by Duncan's multiple-range test……..……..... 46

Sumário

Dedicatória v
Agradecimentos vi
Resumo vii
Abstract viii
Lista de figuras ix
Lista de tabelas x
1. Introdução 12
2. Revisão de literatura 15
3. Referências bibliográficas 20
4. Artigo Científico 26
Effects of stocking density on growth, physiological responses and flesh quality in juvenile cobia (*Rachycentron canadum*) 27
1. Introduction 28
2. Material and methods 30
3. Results 34
4. Discussion 35
5. Acknowledgements 40
6. References 40
5. Anexo (Normas para publicação na Aquaculture) 49

1. Introdução

Nas últimas três décadas, a aquicultura mundial tem apresentado uma taxa de crescimento anual de 8,8%, sendo a atividade que mais cresce dentre os setores de produção animal (FAO, 2012a). Com o crescimento da indústria, produtores seguem uma tendência de maximizar a densidade de estocagem dos animais, visando o aumento nos níveis de produção. No entanto, altas densidades de estocagem podem provocar estresse, comprometendo não apenas o crescimento, mas também a reprodução, imunidade e bem estar dos peixes em cativeiro (ROWLAND et al., 2006; OBA et al., 2009), refletindo negativamente na produtividade e consequentemente na economia de empreendimentos aquícolas.

O beijupirá (*Rachycentron canadum*) é uma espécie emergente no cenário da aquicultura mundial, sendo apontado como um candidato à piscicultura marinha no Brasil (CAVALLI e HAMILTON, 2009). Em 2010, a produção mundial superou 40 mil toneladas (FAO, 2012b). Entre os principais países produtores dessa espécie se destacam a China e Taiwan. No Brasil, alguns estados já apresentam iniciativas de produção (CAVALLI et al., 2011; SAMPAIO et al., 2011). O beijupirá é comumente criado em duas fases. A primeira é realizada no continente, com as etapas de reprodução e larvicultura, enquanto que a segunda fase, denominada engorda, geralmente é realizada em gaiolas flutuantes instaladas em águas costeiras ou em mar aberto. Consequentemente, as opções de tratamento de doenças na fase de engorda são bastante restritas devido a questões logísticas e práticas, considerando principalmente a localização e eventualmente o tamanho das estruturas onde os peixes são criados. Por outro lado, a prevenção é a estratégia mais viável para o manejo de doenças na criação de beijupirá, e deve ser realizada de maneira a evitar a exposição dos peixes a situações estressantes (TRUSHENSKI et al., 2010). Deste modo, se torna imprescindível determinar a densidade de estocagem ótima nas fases iniciais da criação como forma de minimizar o estresse e, consequentemente, otimizar a produção.

Ashley (2007) comenta que a determinação da densidade ótima deve encontrar um ponto de equilíbrio entre a máxima produtividade e o mínimo de fatores que causem desordens fisiológicas e comportamentais nos peixes. Utilizando sistemas de recirculação de água, Liao et al. (2004) notaram que a densidade ótima para o beijupirá com peso entre 4 e 8 g seria de 370 peixes/m^3 ou cerca de 28 kg/m^3, enquanto Webb et al. (2007) não encontraram diferenças no crescimento quando peixes de quase 7 g foram criados durante 10 semanas em densidades variando de 40 a 440 g/m^3. No entanto, estudos sugerem que juvenis de beijupirá podem ser afetados por estressores associados à alta densidade em sistemas de recirculação (SCHWARZ et al., 2007) ou em gaiolas (BENETTI et al., 2008), em que infecções relacionadas ao estresse e infestações de *Pasteurella*, *Vibrio*, *Amyloodinium*, e *Benedenia spp*. são comuns (CHEN et al., 2001b; LOPEZ et al., 2002; CHI et al., 2003, KERBER et al., 2011).

A resposta primária ao estresse envolve o aumento de catecolaminas e cortisol (BARTON e IWAMA, 1991; BARTON, 2002). Esses hormônios induzem respostas secundárias que são caracterizadas por uma redução no glicogênio hepático, aumento nos níveis de glicose e lactato plasmáticos, assim como disfunção na osmoregulação. Portanto, o estresse pode levar a um maior consumo das reservas energéticas e esta relocação de energia metabólica pode interferir negativamente em outros processos fisiológicos, como crescimento, reprodução e imunidade (BARTON e IWAMA, 1991; WENDELAAR BONGA, 1997; MOMMSEN et al. 1999).

A literatura especializada ainda conta com poucos estudos tratando da resposta fisiológica do beijupirá ao estresse. Dois estudos (CNAANI e McLEAN, 2009; TRUSHENSKI et al., 2010) avaliaram a resposta de juvenis expostos a estressores experimentais agudos, como exposição ao ar ou ao baixo volume de água, e indicaram que os níveis de glicose, osmolalidade, cortisol e lactato no sangue aumentaram nas primeiras horas após a exposição aos estressores, retornando aos níveis normais em até 12 h.

Diversas espécies têm sido amplamente estudadas quanto às influências do estresse sobre a qualidade de filé, no entanto nenhuma informação se encontra disponível na literatura especializada a respeito do beijupirá. A maioria dos estudos tem abordado os efeitos de estressores agudos, que comumente podem ocorrer durante a despesca ou métodos de abate. Dois indicadores frequentemente utilizados para avalição da qualidade de filé são a perda de líquido por exsudação e o rigor mortis. Em diversas espécies, o aumento da atividade e ocorrência de estresse antes do abate, somado as subsequentes respostas endócrinas, podem resultar em uma rápida queda do pH muscular devido ao aumento de ácido lático oriundo do metabolismo anaeróbio do músculo. A redução do pH muscular resulta em um rápido início do rigor mortis, podendo causar uma redução na vida de prateleira e aumento na perda de liquido por exsudação (BERG et al., 1997; JERRETT et al., 1996; POLI et al., 2005).

A perda de líquido por exsudação, por sua vez, se refere a perdas de fluídos de conteúdo proteico do interior das células, denominado exsudato, podendo causar uma degeneração mais rápida do tecido muscular. Dessa forma, uma grande quantidade de exsudato pode influenciar a aparência, suculência, textura e sabor do filé, a qual, portanto, representa um fator de impacto comercial (KRISTOFFERSEN et al., 2007). Por outro lado, o rigor mortis é descrito como o primeiro processo post mortem que apresenta uma maior influencia sobre aparência e textura do filé. O início e desenvolvimento do rigor mortis são afetados por diversos fatores, como espécie, idade e tamanho dos peixes, procedimentos pré-abate e os métodos de abate (BERG et al., 1997; VAN DE VIS et al., 2003; POLI et al., 2005; SIMITZIS et al., 2013).

Dessa forma, o presente estudo avaliou a resposta fisiológica, desempenho zootécnico e qualidade de filé de juvenis de beijupirá submetidos a um estressor crônico, como altas densidades, visando a definição de condições de criação que levem à melhoria do bem estar animal e ao incremento da produtividade e qualidade do produto final.

2. Revisão de literatura

Atualmente, existem várias iniciativas de criação do beijupirá (*R. canadum*) em águas brasileiras, mais especificamente nos estados da Bahia, Espírito Santo, Pernambuco, São Paulo (CAVALLI et al., 2011) e Rio Grande do Norte. Apesar dessas iniciativas, a produção comercial de peixes marinhos no Brasil não tem registros nas estatísticas de produção de pescado (MPA, 2012).

O beijupirá é uma espécie pelágica e migratória de ampla distribuição geográfica, ocorrendo em águas tropicais e subtropicais em todos os continentes entre as latitudes de 32ºN e 28ºS, com exceção da porção leste do Oceano Pacífico (SHAFFER e NAKAMURA, 2006). Devido à ausência de uma vesícula gasosa, possui hábito natatório ativo similar ao observado nos tubarões de pequeno porte. Possui hábito alimentar predador, composto por peixes, crustáceos e, eventualmente bivalves (ARENDT et al., 2001) e lulas (FRANKS et al., 1996), embora no litoral Pernambucano sua alimentação seja composta por peixes ósseos demersais, ingerindo poucos crustáceos (DOMINGUES et al., 2007). O hábito alimentar indica, portanto, uma preferência alimentar carnívora associada à disponibilidade de alimento no ambiente natural ao longo do litoral.

Nos últimos anos, o beijupirá tem sido alvo de uma série de estudos, nos quais foi demonstrado ser um excelente candidato para a criação. Entre as principais características de interesse estão a alta taxa de crescimento (ARNOLD et al., 2002; LIAO et al., 2004), eficiência alimentar e bom valor comercial (CHOU et al., 2001; WANG et al., 2005), excelente qualidade de carne, ideal para a preparação de sashimi (CHEN, 2001[a]; CHOU et al., 2001; KAISER e HOLT, 2005), adaptação ao confinamento (SUN et al., 2006), relativa tolerância à baixas salinidades (FAULK e HOLT, 2006; RESLEY et al., 2006) e facilidade em desovar em cativeiro (ARNOLD et al., 2002; FAULK e HOLT, 2006).

Devido a estas características, diversos países têm demonstrado interesse em produzi-lo em escala comercial, sendo China e Taiwan os países que lideram a produção mundial, onde a criação ocorre de forma intensiva desde a década de 1990 (LIAO et al., 2004). Outros países também já

figuram na lista de produtores de beijupirá, entre os quais Porto Rico e Vietnã (BENETTI, et al., 2007; NHU et al., 2011). Portanto, a maioria dos resultados positivos com a criação do beijupirá está concentrada no sudeste asiático, sendo que em Taiwan cerca de 80% das gaiolas são destinadas a criação do beijupirá (LIAO et al., 2004). É importante salientar que algumas regiões produtoras do sudeste asiático possuem um inverno rigoroso quando comparado ao Brasil. Segundo Yu e Ueng (2007), o beijupirá é criado em regiões com variação anual de temperatura de 15,5 a 30,5ºC, sendo o seu crescimento acelerado em temperaturas acima de 28ºC.

Em sistemas de criação em alto mar, o beijupirá pode crescer de 4 a 6 kg por ano (ARNOLD et al., 2002; CHOU et al., 2001; WANG et al., 2005) e até 10 quilos em um período de 12 a 14 meses (LIAO et al., 2004). Em Porto Rico, estudos mostraram que o peso final pode variar de 1,5 a 6 kg em 12 meses (BENETTI et al., 2007). O tamanho comercial no mercado mundial é de 6 a 8 kg, enquanto que no mercado interno de Taiwan há uma preferência por peixes variando entre 8 e 10 kg (MIAO et al., 2009).

A dieta utilizada é nutricionalmente balanceada, normalmente possuindo 45% de proteína bruta, de 15 a 16% de lipídio bruto e 11% de cinzas (CHOU et al., 2001; LIAO et al., 2004). No entanto, as dietas podem ser manipuladas para se obter uma maior concentração de lipídios na carne (GAYLORD e GATLIN, 2000), sem prejuízo na conversão alimentar aparente do beijupirá (CHOU et al., 2001). Apesar da exigência mínima de lipídio do beijupirá ser de 6% (CHOU et al., 2001), é possível utilizar uma dieta com maiores porcentagens de lipídio bruto (CRAIG et al., 2006). A maior vantagem do enriquecimento das dietas com lipídios está relacionada à presença dos ácidos graxos altamente insaturados (HUFAs), principalmente o EPA (ácido eicosapentanóico) e o DHA (ácido decosapentanóico), que juntos são responsáveis pela boa qualidade nutricional das dietas (WATANABE, 2002). O mercado japonês, por exemplo, exige um produto final com alta qualidade para seu consumo *in natura* na forma de sashimi (CRAIG et al., 2006), portanto sendo necessário uma suplementação de lipídios na dieta superior ao exigido pela espécie.

O fator de conversão alimentar (FCA) é relativamente baixo, variando entre 1,02 e 1,80, dependendo do tamanho dos peixes (SU et al., 2000), das condições do ambiente, principalmente a temperatura, e proporção de farinha de peixe utilizada nas dietas. Entretanto, normalmente são obtidos valores de FCA em torno de 2,0, com sobrevivência acima de 75% (BENETTI et al., 2007).

Na fase de berçário, as criações comerciais de beijupirá em Taiwan utilizam normalmente densidade de 13 peixes/m^3 (≈400 g/m^3), alcançando uma produtividade final de 8,4 kg/m^3 em um período que varia entre 4 e 5 meses de criação. Já na fase de engorda final, os peixes são transferidos para gaiolas maiores onde são criados com taxas de densidade de 2 peixes/m^3, alcançando uma produtividade de 14 kg/m^3 ao final de 6 a 8 meses de criação (LIAO et al., 2004). Em Porto Rico, a produtividade final numa criação em mar aberto variou entre 5 e 15 kg/m^3, mas o ganho de peso e a sobrevivência diminuíram nas densidades mais altas (BENETTI et al., 2010). A variabilidade na produtividade indica uma relação direta da densidade de estocagem inicial com o ganho de peso, mortalidade e conversão alimentar.

A densidade de estocagem é amplamente reconhecida como um fator crítico na aquicultura, uma vez que altas densidades podem representar uma potencial fonte de estresse nos peixes. A resposta primária ao estresse envolve o aumento de catecolaminas e cortisol (BARTON e IWAMA, 1991; BARTON, 2002). Esses hormônios induzem respostas secundárias que são caracterizadas por uma redução no glicogênio hepático, aumento nos níveis de glicose e lactato plasmáticos, assim como disfunção na osmoregulação. Portanto, o estresse pode levar a um maior consumo das reservas energéticas e esta relocação de energia metabólica pode interferir negativamente em outros processos fisiológicos, como crescimento, reprodução e imunidade (BARTON e IWAMA, 1991; WENDELAAR BONGA, 1997; MOMMSEN et al. 1999).

A literatura especializada ainda conta com poucos estudos tratando da resposta fisiológica do beijupirá ao estresse. Dois estudos (CNAANI e McLEAN, 2009; TRUSHENSKI et al., 2010) avaliaram a resposta de juvenis expostos a estressores experimentais agudos, como exposição ao ar

ou ao baixo volume de água, e indicaram que os níveis de glicose, osmolalidade, cortisol e lactato no sangue aumentaram nas primeiras horas após a exposição aos estressores, retornando aos níveis normais em até 12 h.

Espécies como salmão do Atlântico, robalo europeu e pargo europeu são amplamente estudadas quanto as influencias do estresse sobre a qualidade de filé (ROTH et al., 2006; BAGNI et al., 2007; MØRKØRE et al., 2008), mas no entanto nenhuma informação se encontra disponível na literatura especializada a respeito do beijupirá. A maioria dos estudos tem abordado os efeitos de estressores agudos, que comumente podem ocorrer durante a despesca ou métodos de abate. Por exemplo, uma exaustiva atividade muscular antes do abate pode provocar uma drástica redução de ATP muscular e produção de ácido lático, este último resultante do excessivo consumo de glicogênio, e que provoca uma concomitante redução do pH muscular (BERG et al., 1997; JERRETT et al., 1996; POLI et al., 2005). Portanto, o aumento da atividade e o estresse antes do abate pode antecipar o desenvolvimento do rigor mortis, reduzir a vida de prateleira e aumentar a perda de liquido por exsudação.

Dessa forma, dois indicadores frequentemente utilizados para avalição da qualidade de filé são a perda de líquido por exsudação e o rigor mortis. A perda de líquido por exsudação se refere a perdas de fluídos de conteúdo proteico do interior das células, denominado exsudato, podendo causar uma degeneração mais rápida do tecido muscular. Dessa forma, uma grande quantidade de exsudato pode influenciar a aparência, suculência, textura e sabor do filé, a qual, portanto, representa um fator de importância comercial (KRISTOFFERSEN et al., 2007). Por outro lado, o rigor mortis é descrito como o primeiro processo post mortem que apresenta uma maior influencia sobre aparência e textura do filé. O início e desenvolvimento do rigor mortis são afetados por diversos fatores, como espécie, idade e tamanho dos peixes, procedimentos pré-abate e os métodos de abate (BERG et al., 1997; VAN DE VIS et al., 2003; POLI et al., 2005; BAGNI et al., 2007; SIMITZIS et al., 2013).

Nenhuma informação está disponível na literatura especializada a respeito dos efeitos de estressores crônicos sobre as respostas fisiológicas e qualidade do filé do beijupirá. Dessa forma, o presente estudo investigou a resposta fisiológica, desempenho zootécnico e qualidade de filé de juvenis de beijupirá submetidos a diferentes densidades, visando a definição de condições de criação que levem à melhoria do bem estar animal e ao incremento da produtividade e qualidade do produto final.

3. Referências bibliográficas

ARENDT, M.D.; OLNEY, J.E.; LUCY, J.A. Stomach content analysis of cobia, *Rachycentron canadum*, from lower Chesapeake bay. **Fishery Bulletin,** v.99, p.665-670, 2001.

ARNOLD, C. R.; KAISER, J. B.; HOLT, G. J. Spawning of cobia (*Rachycentron canadum*) in captivity. **Journal of the World Aquaculture Society**, v. 33, n.2, p. 205-208, 2002.

ASHLEY, P. J. Fish welfare: Current issues in aquaculture. **Applied Animal Behaviour Science**, v. 104, p. 199-235, 2007.

BAGNI, M., CIVITAREALE, C., PRIORI, A., BALLERINI, A., FINOLA, M., BRAMBILLA, G., MARINO, G., 2007. Pre-slaughter crowding stress and killing procedures affecting quality and welfare in sea bass (*Dicentrarchus labrax*) and sea bream (*Sparus aurata*). **Aquaculture** 263, 52–60.

BARTON B.A. Stress in fishes: a diversity of responses with particular reference to changes in circulating costicosteroids. **Integrative and Comparative Biology**, v. 42, p. 517-525, 2002.

BARTON B.A.; IWAMA G.K. Physiological changes in fish from stress in aquaculture with emphasis on the response and effects of corticosteriods. **Annual Review of Fish Diseases**, v. 129, p. 3-26, 1991.

BENETTI, D.; ORHUN, R.; ZINK I; CAVALIN, F.G.; SARDENBERG B.; PALMER, K.; DENLINGER, B.; BACOAT D.; O'HANLON, B. Aquaculture of cobia (*Rachycentron canadum*) in the Americas and the Caribbean. In: I.-C. LIAO (ed), Cobia Aquaculture, p.20, 2007.

BENETTI, D.D., ORHUM, M.R., SARDENBERG, B., O'HALON, B., WELCH, A., HOENIG, R., ZINK, I., RIVERA, J.A., DENLINGER, B., BACOAT, D., PALMER, K., CAVALIN, F. Advances in hatchery and grow-out technology of cobia *Rachycentron canadum* (Linnaeus). **Aquaculture** Research, v. 39, p. 701–711, 2008.

BENETTI, D.D.; O'HANLON, B.; RIVIERA, J.A.; WELCH, A.W.; MAXEY, C.; ORHUN, M.R. Growth rates of cobia **(***Rachycentron canadum***)** cultured in open ocean submerged cages in the Caribbean. Aquaculture, v. 302, p. 195-201, 2010.

BERG, T., ERIKSON, U., NORDVEDT, T.S. Rigor mortis assessment of Atlantic salmon (*Salmo salar*) and effects of stress. **Journal of Food Science,** v. 62, p. 439–446, 1997.

CAVALLI, R. O.; HAMILTON, S. Piscicultura marinha no Brasil com ênfase na produção do beijupirá. **Revista Brasileira de Reprodução Animal**, v. 6, p. 64-69, 2009.

CAVALLI, R.O., DOMINGUES, E.C., HAMILTON, S. Desenvolvimento da produção de peixes em mar aberto: possibilidades e desafios. **Revista Brasileira de Zootecnia**, v. 40, p.155-164, 2011.

CHEN, B.S. Studies on the net-cage culture and diseases control technology of cobia, *Rachycentron canadum* (Linnaeus). **6th Asian Fisheries Book of Abstracts**. Asian Fisheries Society, Manila, Philippines, 2001a.

CHEN, S.-C., KOU, R.-J., WU, C.-T., WANG, P.-C., SU, F.-Z. Mass mortality associated with a *Sphaerospora*-like myxosporidean infestation in juvenile cobia, *Rachycentron canadum* (L.), marine cage cultured in Taiwan. **Journal of Fish Diseases**, v. 24, p. 189–195, 2001b.

CHI, S.C., SHIEH, J.R., LIN, S.J. Genetic and antigenic analysis of betanodaviruses isolated from aquatic organisms in Taiwan. **Diseases of Aquatic Organisms**, v. 55, p. 221–228, 2003.

CHOU, R.-L., SU, M.S., CHEN, H.Y. Optimal dietary protein and lipid levels for juvenile cobia (*Rachycentron canadum*). **Aquaculture**, v.193, p.81-89, 2001.

CNAANI, A., MCLEAN, E. Time-course response of cobia (*Rachycentron canadum*) to acute stress. **Aquaculture**, v. 289, p. 140–142, 2009.

CRAIG, S.R., SCHWARZ, M.H., MCLEAN, E. Juvenile cobia (*Rachycentron canadum*) can utilize a wide range of protein and lipid levels without impacts on production characteristics. **Aquaculture**, v. 261, p. 384–391, 2006.

DOMINGUES, E. C., PEREGRINO JR., R. B., MANZELLA JR., J. C., VASKE JR.,T., HAZIN, F.H.V., CAVALLI, R. O., SEVERI, W., HAMILTON, S. 2007. Aspectos biológicos do beijupirá, *Rachycentron canadum*, espécie potencial para o desenvolvimento da piscicultura marinha no Nordeste. In: II Seminário de Piscicultura Alagoana, 2007. Penedo. Anais. Penedo/AL. SEBRAE-AL. Brasil.

FAO. Food and Agriculture Organization of the United Nations. The state of world fisheries and aquaculture 2012. Rome, 2012a.

FAO. Cultured Aquatic Species Information Programme- *Rachycentron canadum*. Food and Agricultural Organization, Rome, 2012. http://www.fao.org/fishery/culturedspecies/ Rachycentron_canadum/en Acesso em 06/14/2012b.

FAULK, C.K. & HOLT, G.J. Responses of cobia *Rachycentron canadum* larvae to abrupt or gradual changes in salinity. **Aquaculture**, v.254, p.275-283, 2006.

FRANKS, J.S.; GARBER, N.M.; WARREN, J.R. Stomach contents of juvenile cobia, *Rachycentron canadum*, from the Northern Gulf of Mexico. **Fishery Bulletin**, v. 94, p. 374–380, 1996.

GAYLORD, T.G., GATLIN III, D.M. Dietary lipid level but not L-carnitine affects growth performance of hybrid striped bass (*Morone chrysops* ♀ x *M. saxatilis* ♂). **Aquaculture**, v. 190, p. 237–246, 2000.

KAISER, J.B., G.J. HOLT. Species Profile Cobia. Southern Regional Aquaculture Center Publication, n.7202, p.6, 2005.

KERBER, C.E.; SANCHES, E.G.; SANTIAGO, M.; LUQUE, J. First record of *Neobenedenia melleni* in sea-farmed cobia (*Rachycentron canadum*) in Brazil. **Rev. Bras. Parasitol. Vet.**, v. 20, n.4, 331-333, 2011.

KRISTOFFERSEN, S., VANG, B., LARSEN, R., AND OLSEN, R. L.. Pre-rigor filleting and drip loss from fillets of farmed Atlantic cod (*Gadus morhua* L.). **Aquaculture Research**, v. 38, p. 1721-1731, 2007.

LIAO, I.C.; HUANG, T.S.; TSAI, W,-S.; HSUEH, C.M.; CHANG, S.L.; LEAÑO, E.M. Cobia culture in Taiwan: current status and problems. **Aquacultur**e, v. 237, p. 155-165, 2004.

LOPEZ, C., RAJAN, P.R., LIN, J.H.-Y., KUO, T.-Y., YANG, H.-L. Disease outbreak in seafarmed cobia (*Rachycentron canadum*) associated with *Vibrio spp*., *Photobacterium damselae spp. piscida*, *monogenean* and *myxoxporean* parasites. **Bulletin of the European Association of Fish Pathologists**, v. 22, p. 206–211, 2002.

MIAO, S.; JEN, C. C.; HUANG, C. T.; HU, S-H. Ecological and economic analysis for cobia, *Rachycentron canadum*, commercial cage culture in Taiwan. **Aquaculture International**, v.17, p.125-141, 2009.

MOMMSEN, T.P., VIJAYAN, M.M., MOON, T.W. Cortisol in teleosts: dynamics, mechanisms of action, and metabolic regulation. **Reviews in Fish Biology and Fisheries**, v. 9, p. 211–268, 1999.

MPA - Ministério da Pesca e Aquicultura. **Boletim estatístico da Pesca e Aquicultura – Brasil 2010**. Brasília, DF, Brasil: Ministério da Pesca e Aquicultura, 2012. 128pp. 2012.

NHU, V.C.; NGUYEN, H.Q.; LE, T.L.; TRAN, M.T.; SORGELOOS, P.; DIERCKENS, K.; REINERTSEN, H.; KJORSVIK, E.; SVENNEVIG, N. Cobia *Rachycentron canadum* aquaculture in Vietnam: Recent developments and prospects. **Aquaculture**, v. 315, p. 20-25, 2011.

OBA, E. T.; MARIANO, W. S.; SANTOS, L. R. B. Estresse em peixes cultivados: agravantes e atenuantes para o manejo sustentável. In: TAVARES-DIAS, M. (Org). 2009. Manejo e Sanidade de Peixes em Cultivo. Macapá: Empraba Amapá. p. 389-424, 2009.

POLI, B.M., PARISI, G., SCAPPINI, F., ZAMPACAVALLO, G. Fish welfare and quality as affected by pre-slaughter and slaughter management. **Aquaculture International**, v. 13, p. 29–49, 2005.

RESLEY, M. J.; WEBB, K. A.; HOLT, G. J. Growth and survival of juvenile cobia, *Rachycentron canadum*, at different salinities in a recirculating aquaculture system. Aquaculture, v. 253, p. 398-407, 2006.

ROWLAND, S. J.; MIFSUD, C.; NIXON, M.; BOYD, P. Effects of stocking density on the performance of the Australian freshwater silver perch (*Bidyanus bidyanus*) in cages. **Aquaculture**, v. 253, p. 301-308, 2006.

SAMPAIO, L.A., MOREIRA, C.B., MIRANDA-FILHO, K.C., ROMBENSO, A.N. Culture of cobia *Rachycentron canadum* in near-shore cages off the Brazilian coast. **Aquaculture Research**, v. 42, p. 832-834, 2011.

SCHWARZ, M.H., MCLEAN, E., CRAIG, S.R. Research experience with cobia: larval rearing, juvenile nutrition and general physiology, in: Liao, I.C., Leaño, E.M. (Eds.), Cobia Aquaculture: Research, Developments And Commercial Production. Co-Published by: Asian Fisheries Society; The Fisheries Society of Taiwan; World Aquaculture Society; and National Taiwan Ocean University, Keelung, pp. 1–1, 2007.

SHAFFER, R. V.; NAKAMURA, E. L. Synopsis of biological data on the cobia *Rachycentron canadum* (Pisces: Rachycentridae). U.S. Department of Commerce, NOAA Technical Report. Washington D.C. FAO Fisheries Synopsis, n.153, 2006.

SIMITZIS, P.E., TSOPELAKOS, A., CHARISMIADOU, M.A., BATZINA, A., DELIGEORGIS, S.G., MILIOU, H., 2013. Comparison of the effects of six stunning/killing procedures on flesh quality of sea bass (*Dicentrarchus labrax*, Linnaeus 1758) and evaluation of clove oil anaesthesia followed by chilling on ice/water slurry for potential implementation in aquaculture. **Aquaculture Research** *in press*, 1-12. doi: 10.1111/are.12120

SU, M.S.; CHIEN, Y.H.; LIAO, I.C. Potential of marine cage aquaculture in Taiwan: cobia culture. In: Liao, I.C., Lin, C.K. (Eds.), Cage Aquaculture in Asia: Proceedings of the First International Symposium on Cage Aquaculture in Asia (Ed. LIAO, I.C.; LIN, C.K.), pp. 97-106. Asian Fisheries Society, Manila, and World Aquaculture Society- Southeast Asian Chapter, Bangkok. 2000.

SUN, L., CHEN, H., HUANG, L., WANG, Z.,YAN, Y. Growth and energy budget of juvenile cobia (*Rachycentron canadum*) relative to ration. **Aquaculture**, v. 257, p. 214-220, 2006.

TRUSHENSKI. J., SCHWARZ, M., TAKEUCHI, R., DELBOS B., SAMPAIO, L.A. Physiological responses of cobia *Rachycentron canadum* following exposure to low water and air exposure stress challenges. **Aquaculture**, v. 307, p. 173-177, 2010.

VAN DE VIS, H., KESTIN, S., ROBB, D., OEHLENSCHLAGER, J., LAMBOOIJ, B., MUNKNER, W., KUHLMANN, H., KLOOSTERBOER, K., TEJADA, M., HUIDOBRO, A., OTTERA, H., ROTH, B., SØRENSEN, N.K., AKSE, L., BYRNE, H. AND NESVADBA, P., 2003. Is humane slaughter of fish possible for industry? **Aquaculture Research**, v. 34, p. 211–220, 2003.

WANG, J-T.; LIU, Y-J.; TIAN, L-X.; MAI, K-S.; DU, Z-Y.; WANG, Y.; YANG, H-J. Effect of dietary lipid level on growth performance, lipid deposition, hepatic lipogenesis in juvenile cobia (*Rachycentron canadum*). **Aquaculture**, v. 249, p. 439-447, 2005.

WATANABE, T. Strategies for further development of aquatic feeds. **Fisheries Science**, v.68, n.2, p.242-252, 2002.

WEBB, K.A.; HITZFELDER G.M.; FAULK, C.K.; HOLT. G.J. Growth of juvenile cobia, *Rachycentron canadum*, at three different densities in a recirculating aquaculture system. **Aquaculture**, v. 264, p. 223–227, 2007.

WENDELAAR BONGA S.E. The stress response in fish. **Physiological Reviews**. v. 7, p. 591-625, 1997.

YU, S. L.; UENG, P. S. Impact of water temperature on growth in cobia, *Rachycentron canadum*, cultured in cages. **Israeli Journal of Aquaculture-Bamidgeh**, v.59, p.47-51, 2007.

4. Artigo Científico

Artigo científico a ser encaminhado a Revista: Aquaculture

Todas as normas de redação e citação, deste capítulo, atendem as estabelecidas pela referida revista (em anexo).

Effects of stocking density on growth, physiological responses and flesh quality in juvenile cobia (*Rachycentron canadum*)

Victor A. Silva[1,2]*, Jesse Trushenski[3], Michael H. Schwarz[4], Ronaldo O. Cavalli[1]

[1] Federal Rural University of Pernambuco – UFRPE, Department of Fisheries and Aquaculture, Laboratory of Marine Fish Farming, Av. Dom Manuel de Medeiros, s/n, Dois Irmãos, 52171-900, Recife, PE, Brazil

[2] International Initiative for Sustainable and Biosecure Aquafarming, Norfolk, VA 23053, USA

[3] Fisheries and Illinois Aquaculture Center, Southern Illinois University Carbondale, Carbondale, Illinois, 62901, USA

[4]Virginia Seafood Agricultural Research and Extension Center, Virginia Tech, 102 South King Street, Hampton, Virginia, 23669, USA

* Corresponding author

Phone: +55 (81) 3320.6524, Fax: +55 (81) 3320.6502

E-mail: s.victorandrade@gmail.com

Abstract

Stocking density is recognized as a critical factor in aquaculture, since high densities represent a potential source of chronic stress. This study investigated the effects of stocking density on growth, physiological responses and flesh quality of juvenile cobia (*Rachycentron canadum*). The trial was conducted for 42 days in a recirculating aquaculture system equipped with 70-L tanks. Cobia of 46.32 ± 0.22 g (mean ± SEM) were randomly distributed into three stocking densities (3.30 ± 0.02; 6.67 ± 0.03; 13.15 ± 0.16 g L^{-1}), and three replicates. Treatment groups were nominally assigned as low density (LD; 5 fish/tank), medium density (MD; 10 fish/tank), and high density (HD; 20 fish/tank), respectively. Plasma concentrations of cortisol and lactate, and osmolality were not

affected by density levels, and values were according with those reported as basal for juvenile cobia. Weight gain and specific growth rate were significantly higher in the LD group. Drip loss had no significant difference among density groups, but stocking density influenced the onset and development of rigor mortis. Our findings indicated that rearing juvenile cobia at the high densities negatively affected growth rate and flesh quality. Juvenile cobia reared at the high density were able to cope with stress in terms of plasma composition, but energy from the diet and/or body reserves were probably diverted from growth as a response to the elevated metabolic demand caused by stress. Juvenile cobia should be maintained at a final biomass of up to approximately 15 kg m^{-3} to ensure a maximal growth rate and an improved final product quality. To our knowledge, this is the first report assessing the influence of a chronic stressor on both physiological responses and flesh quality parameters in cobia.

Keywords: Fish, chronic stress, hematology, rigor mortis, drip loss.

1. Introduction

In the last three decades, aquaculture has expanded at an average annual rate of 8.8% (FAO, 2012). As the aquaculture industry grows, farmers tend to increase fish stocking densities in order to achieve higher production levels. However, increased stocking densities may induce stress, and lead not only to reduced survival and impaired growth rates, reproduction and immune status, but ultimately affect the profitability of the activity.

High stocking densities in fish grown in captivity represents a potential source of chronic stress, and it is widely recognized as a critical factor in aquaculture. Increases in the concentrations of plasma catecholamines and cortisol are described as primary responses to stress in fish. These hormones lead to secondary responses, which involve metabolic energy demand to cope with stress. Decrease in hepatic glycogen, and increase in concentrations of plasma glucose and lactate, as well as a temporary dysfunction in osmoregulation are described as secondary stress responses. Thus, stress may induce energy consumption, which negatively interferes on growth, reproduction and

immunity (Barton & Iwama, 1991; Wendelaar Bonga, 1997; Mommsen et al., 1999; Barton, 2002). Therefore, the recognition of stress status is critical to ensure fish welfare and the success of any aquaculture operation. Many quantitative parameters, such as growth rate, condition factor, hepatosomatic index and blood composition (concentration of plasma cortisol, osmolality and lactic acid), may be used as indicators of stress in fish (Barton & Iwama, 1991; Montero et al., 1999).

Cobia (*Rachycentron canadum*) is an emerging species for aquaculture. In 2010, world production was estimated at around 40,000 MT (FAO, 2013). The main producing areas are in China and Taiwan, but several other countries in the Americas and Asia have demonstrated interest in cobia aquaculture. Commercial culture is basically divided in two major steps. The first one occurs in land-based rearing systems, where early juveniles are produced. In the second step, known as grow-out, fish are usually reared in cages placed in coastal or open sea waters. Under these conditions, disease treatment options are restricted due to practical and logistic limitations. As a result, strategies to avoid or minimize stress exposure and hence reduce the occurrence of diseases and parasites are an important management strategy.

Determining the optimum fish stocking density is one way to diminish stress. In this regard, it is essential to establish a balance between increasing productivity and minimizing factors that may lead to physiologic and behavior disorders (Conte, 2004; Ashley, 2007). Liao et al. (2004), Webb et al. (2007) and Benetti et al. (2010) evaluated the influence of stocking density on growth performance, but none of them considered the effects on the physiological responses of cobia. Currently, only two studies considering the effects of acute stress on cobia are available. Cnaani & McLean (2009) assessed the responses of juvenile cobia exposed to an experimental stressor (air exposure) over a period of 24 hours, while Trushenski et al. (2010) evaluated the hematological responses of cobia exposed to low water and air exposure over a 72-hour time frame.

While the effects of stress on flesh quality have been assessed in several fish species, again no information is available for cobia. Most studies have addressed the effects of acute stressors during pre-slaughter procedures or slaughtering methods. Exhaustive exercise and occurrence of

severe stress during slaughter may result in a rapid drop in muscle pH due to the production of lactic acid derived from anaerobic metabolism of white muscle. Reduced muscle pH results in a rapid onset of rigor mortis, which may reduce shelf life, alter flesh texture and increase drip loss (Berg et al., 1997; Jerrett et al., 1996; Poli et al., 2005). Drip loss and rigor mortis are therefore two commonly used indicators of flesh quality. Accordingly, this study evaluated the growth performance, physiological responses and flesh quality of juvenile cobia exposed to a chronic stressor such as high stocking density.

2. Material and methods

The study was performed at the Virginia Tech-Virginia Seafood Agricultural Research and Extension Center (VSAREC) in Hampton, VA, USA. The trial was conducted for 42 days in a RAS containing 70-L tanks, fluidized-bed biofilters, bubble bead filters, UV sterilizers, immersion titanium heaters, protein skimmers for processing/removal of dissolved organics and small suspended solids, and a diffusion aeration system. Fish were fed a commercial carnivorous fish diet (Otohime, Marubeni Nisshin Feed Company, Japan; 48% protein, 13% lipids, 4 mm) twice daily to apparent satiation. Water temperature, salinity, and dissolved oxygen (DO) were monitored daily (YSI-85 series dissolved oxygen meter; YSI Inc., Yellow Springs, Ohio, USA). Total ammonia-, nitrite- and nitrate-nitrogen (Spectrophotometric analysis; Hach Inc., Loveland, Colorado, USA), pH (YSI pH100 meter; YSI Inc., Yellow Springs, Ohio, USA), and alkalinity (Bromocresol green methyl red titration method, Hach Inc., Loveland, Colorado, USA) were also quantified daily. All fish in this trial were obtained from the same spawn (F3 generation) to avoid eventual biological bias. Prior to stocking the system and beginning the trial, all fish had been fed the same commercial diet that was used throughout the trial.

Tank inflow rates were maintained constant at 12 L min^{-1}, photoperiod was kept on a 12:12 h light/dark cycle, and water quality conditions were maintained as follows (mean ± SEM): temperature = 26.11 ± 0.17 ºC; salinity = 17.6 ± 0.1 g L^{-1}; pH = 7.70 ± 0.03; dissolved oxygen =

6.17 ± 0.07 mg L^{-1}; total ammonia nitrogen = 0.73 ± 0.04 mg L^{-1}; nitrite-nitrogen = 0.31 ± 0.03 mg L^{-1}; nitrate-nitrogen = 35.56 ± 2.30 mg L^{-1} and alkalinity = 142.5 ± 3.20 mg L^{-1}.

Juvenile cobia of 46.32 ± 0.22 g (mean ± SEM) were randomly distributed into three stocking densities (3.30 ± 0.02, 6.67 ± 0.03, 13.15 ± 0.16 g L^{-1}; mean ± SEM), and tested in three replicates. Density groups were nominally assigned as low density (LD; 5 fish/tank), medium density (MD; 10 fish/tank), and high density (HD; 20 fish/tank), respectively.

2.1 Growth performance parameters

On day 42, all fish were counted and survival estimated. Weight gain (WG), specific growth rate (SGR), feed conversion ratio (FCR), condition factor (K), apparent feed intake (FI) and dress out (%) were estimated with the following formulae:

$$\text{WG (\%)} = 100 \times \frac{\text{average final weight} - \text{average initial weight}}{\text{average initial weight}}$$

$$\text{SGR (\%BW day}^{-1}) = 100 \times \frac{\log_e(\text{average final weight}) - \log_e(\text{average initial weight})}{\text{number of days}}$$

$$\text{FCR} = \frac{\text{average individual feed intake}}{\text{average individual weight gain}}$$

$$\text{K} = 1000 \times \frac{\text{individual weight}}{\text{individual lenght}^3}$$

$$\text{FI (\%BW day}^{-1}) = 100 \times \frac{\text{average individual feed intake}}{\frac{(\text{initial individual weight} \times \text{final individual weight})^{0.5}}{\text{days of feeding}}}$$

$$\text{Dress out (\%)} = 100 \times \frac{\text{skinless fillet weight}}{\text{whole body weight}}$$

2.2 Blood sampling

At the end of trial, feeding was suspended 24 h prior to the collection of blood. In each sampling, three fish per tank were netted in a single pass in order to minimize sampling effort and

additive stress associated with repeat netting. Fish were immediately sedated until they could be safely handled in a 100 mg L^{-1} solution of tricaine methanesulfonate (MS-222, Finquel®; Argent Chemical Laboratories, Inc., Redmond, Washington, USA) in culture water. Blood samples were collected from the caudal vasculature using capillary blood collection (Safe-T-Fill®; Kabe Labortechnik, Germany). After samples were collected, fish were euthanized by single cranial pithing. All blood samples were collected within 5 min of capture and sedation to minimize the possibility of other confounding responses. Whole blood were centrifuged (10 min, 10 000 x *g*, 4ºC) and plasma samples were stored frozen (-80ºC) until shipment for later analysis of cortisol, lactate and osmolality.

After blood sampling, whole liver was dissected and weighed for calculation of hepatossomatic index (HSI), which was calculated following the formula $HSI=(W_{liver}/W_{body})x100$. Whereupon, W_{liver} represents liver weight, and W_{body} represents whole body weight. Subsequently, these fish were filleted and samples were collected from the fillet for drip loss assessment. Frozen plasma were packed in ice and transported to the Fisheries and Illinois Aquaculture Center (FIAC), in Carbondale, Illinois, USA, where samples were stored again at -80ºC prior to hematological and analysis.

2.3 Blood plasma analysis

Plasma samples were analyzed at FIAC, and the methodologies were the same as in Trushenski et al. (2010). Briefly, plasma cortisol concentrations were quantified using an immunoassay (Cortisol EIA, DRG International, Mountainside, NJ, USA) according to the manufacturer's instructions. Plasma osmolality concentrations were determined using a vapor pressure osmometer (Model 5100C; Wescor Incorporated, Logan, UT, USA) and plasma lactate concentrations were measured using human blood testing supplies (Accutrend® lactate meter; Roche, Mannheim, Germany).

2.4 Drip loss

All three fish per tank that had their blood collected were filleted, and a slice of approximately 10 g of muscle was sampled from the epixial part of the fillet, weighed (W_0), marked and wrapped in aluminum foil. Samples were stored at 4ºC for a period of 96 hours. Subsequently, samples were unwrapped and weighed again (W_f), and drip loss (DL) was calculated by the following formula: $DL = [(W_0 - W_f) / W_0] \times 100$.

2.5 Rigor mortis

Two fish from each tank were sampled for rigor mortis assessment. A commercial slaughter method was used. Fish were immersed in ice and water (3:1), and slaughtered by gill cutting. Rigor mortis was measured at 0, 1, 3, 5, 7, 9, 12, 24, 36, 48, 72 and 96 hours post mortem using Cuttingers Method (tail drop). The same fish were used for rigor mortis measurements at each time point. In between measurements, fish were maintained at 4ºC. Rigor index (Ir) was calculated by the following formula: $Ir = [(L0 - Lt) / L0] \times 100$ (Bito et al., 1983). Whereupon, L represents the vertical distance between a table surface and the caudal fin base, when half of the fish fork length is placed on the edge of a table. Therefore, L0 represents measurements at time zero, whereas Lt represents subsequent measurements throughout the period of time described above. In this regard, Ir = 100% indicates full rigor.

2.6 Statistical analysis

Although multiple fish were sampled from each tank, replicate tanks served as the experimental units for all statistical analyses (n=3). All production performance and plasma blood data, rigor mortis and drip loss were analyzed by one-way analysis of variance (ANOVA) using the Statistical Analysis System (version 9.1; SAS Institute, Cary, NC, USA) to determine the significance of difference among density treatment means. Duncan's multiple-range test was then performed. Differences were considered significant at $p<0.05$.

3. Results

3.1 Growth performance and survival

Final weight, FCR, FI, HSI, K, and dress-out were not significantly different among density groups (Table 1). On the other hand, weight gain (P = 0.04) and SGR (P = 0.045) were significantly higher in cobia from the LD group than those from either MD or HD groups. Final fish biomass in the LD, MD and HD treatments increased to 15.04, 25.61 and 53.94 kg m^{-3}, respectively. No mortality was registered for fish from the LD and HD groups; however, two fish from the MD group jumped out of the tanks during the trial. Survival was not significantly different between treatments (Table 1)

3.2 Hematological responses

Mean (± SEM) plasma cortisol concentrations ranged from 23.28 to 83.89 ng mL^{-1}, and no significant differences were detected among density groups (p=0.32). Plasma osmolality (p=0.06) and the concentration of lactate (p=0.92) were not influenced by density levels, and values ranged from 366.78 to 390.33 mOsm kg^{-1} and 0.21 to 0.31 mmol L^{-1}, respectively.

3.3 Drip loss

Mean (± SEM) drip loss (%) ranged from 0.85 (± 0.21) to 1.74 (± 0.32). Although drip loss was not significantly different (p=0.057) among density groups, it seemed to be positively associated with stocking density (Figure 1).

3.4 Rigor mortis

At one hour post mortem, the rigor mortis index was significantly larger (p=0.026) in cobia from the HD group compared to those from the MD and LD groups (Figure 2), which indicates an

earlier onset of rigor in juvenile cobia reared at the high density. Afterwards, no significant differences were detected among density groups within each time point. Nevertheless, the average peak for rigor mortis in fish from the HD group was in the range of 1 to 3 hours post mortem, whereas for cobia from the MD and LD groups it was in the range of 3 to 5 h.

4. Discussion

The increase in the concentrations of plasma cathecolamines and cortisol is the most common quantitative indicator of the primary stress responses in fish (Pickering, 1993; Barton, 2002). In this study, however, stocking density had no effect on the plasma cortisol concentrations in cobia after 42 days. Plasma cortisol concentrations may not increase in chronically stressed fish due to a feedback mechanism that causes a down regulation of the hypothalamic-pituitary-interrenal axis. Previous studies (Pickering 1993; Sumpter, 1997; Procarione et al., 1999) demonstrated that in instances of long term stress cortisol concentrations may eventually return to basal levels despite the continued stress, demonstrating habituation to the stressful event.

Temporary osmoregulatory dysfunction is a secondary effect of stress in fish, and increased osmolality would be expected in cobia reared in hypertonic solutions as well as under stress (Davis, 2006; Burkey et al., 2007, Trushenski et al., 2010). However, we found no differences in the osmolality among density groups, which were within values considered as basal for juveniles of this species (Trushenski et al., 2010). Similarly, plasma osmolality and the concentrations of cortisol and lactate did not differ during the a 14 day-long confinement of gilthead sea bream (*Sparus aurata*) at 6 kg m^{-3} and 26 kg m^{-3}, but differences in their hematology were evident after a subsequent exposure to an acute handling stress, indicating that fish experienced a chronic stress situation (Barton et al., 2005).

Increased plasma lactate concentrations is also a secondary response to stress. Although stocking density and plasma lactate correlated positively in previous studies (van de Nieuwegiessen et al., 2008, Costas et al., 2013), density had no effect on the concentration of lactate in juvenile

cobia, and values were within a range considered as basal for this species (Trushenski et al., 2010). In the present study, plasma lactate concentrations are in line with those found in other fish species maintained at high densities (Barton et al., 2005; Herrera et al., 2009; Santos et al., 2010). However, Costas et al. (2013) have recently shown that a high protein diet may represent a metabolic advantage in stressful situations and may counteract immunosuppression in fish. These authors have found that a high (46%) protein diet decreased plasma lactate concentrations in Senegalese sole (*Solea senegalensis*) reared at high densities as well as induced improvements in its immune status. Thus, the relatively high protein diet used in the present study (48%) may also have influenced the present results.

The response of different fish species to the stress caused by increasing density levels varies according to the duration as well as the potency of the stressing factor (Santos et al., 2010). For example, wedge sole (*Dicologoglossa cuneata*) exposed to different stocking densities for a relatively short period of time (22 days) presented no significant differences in growth performance, yet plasma cortisol and osmolality levels were higher in fish at the highest density (Herrera et al., 2009). In contrast, hematological stress indicators (cortisol and proteinemia) and resistance to infection were not different in the European sea bass (*Dicentrarchus labrax*) reared for 63 days at increased density levels, but SGR and feed intake were lower in fish at the highest density (Sammouth et al., 2009). Santos et al. (2010) added that the evaluation of chronic stress is much more challenging than assessing acute stress. They argued that acute stress is often measured as a transient change in the metabolism in an attempt to counteract the stress event, which is not the case of long-term stressing factors.

Differences in weight gain ($p=0.040$) and SGR ($p=0.045$) in this study may not be overly substantial, but they do indicate that raising cobia juveniles at the high and medium densities impaired growth. Decreases in growth rate at increasing stocking densities were observed for several fish species (Canario et al., 1998; Ellis et al., 2002; Sammouth et al., 2009, Roque d'Orbcastel et al., 2010; Santos et al., 2010; Tolussi et al., 2010), including cobia (Liao et al., 2004;

Benetti et al., 2010). Although these negative effects may be due to poor water quality (Bianchini et al., 1996; Santos et al., 2010) and aggressive behavior (Ellis et al., 2002), this does not seem to be the case here as water quality variables were within levels considered appropriate to the development of cobia and no agonistic behavior was observed throughout the experimental period. Interestingly, final biomass in the LD group was around 15 kg m^{-3}, which is similar to what is usually achieved in commercial cobia operations in sea-cage systems (Liao et al., 2004, 2007; Benetti et al., 2010; Sampaio et al., 2011). As fish biomass in the HD and MD exceeded 15 kg m^{-3} during the experimental period, this suggests that the fish biomass may have reached a threshold after which growth would have been negatively affected.

Previous studies demonstrated that cortisol, lactate and osmolality are suitable indicators of acute stress in juvenile cobia (Cnaani & McLean, 2009; Trushenski et al., 2010). For example, Trushenski et al. (2010) analyzed the hematology of cobia exposed to the air and to low water level, and found that osmolality and the concentrations of cortisol, glucose and lactate increased within the first hour after exposure, but returned to basal levels up to 12 h. In this study, the chronic stress caused by stocking density had no effect in these indicators, and their concentrations were within values considered as basal for juvenile cobia. Therefore, if only the results of plasma indicators are considered, one may be led to the conclusion that juvenile cobia experienced no stress. On the other hand, the evaluation of growth performance indicates the opposite.

Studies with sea bass (*D. labrax*) demonstrated that stocking density had no effect on the concentration of plasma stress indicators at the end of the rearing period, but others stress parameters were affected (Di Marco et al., 2008; Santos et al., 2010). A possible explanation to this is the habituation to the stressful conditions by the reduction of physiological alterations during the time of exposure (Pickering 1993; Sumpter, 1997; Procarione et al., 1999; Jentoft et al., 2005; Basrur et al., 2010). The habituation of the stress response, however, does not exclude the effects on growth performance observed in the repeatedly stressed fish versus the unstressed fish (Jentoft et al., 2005). Accordingly, juvenile cobia from this study were able to cope with the chronic stressor,

but this may have caused an important burden in its energy metabolism, since significant differences in growth performance were registered. Stress is an energy-demanding process to which fish usually respond through the mobilization of energy substrates, and this reallocation of metabolic energy negatively influences other physiological process, such as growth, reproduction and immunity (Barton & Iwama, 1991; Iwama, 1998; Mommsen et al., 1999). In *D. labrax (*Di Marco et al., 2008; Santos et al., 2010) and piabanha *Brycon insignis* (Tolussi et al., 2010), increased density levels impaired not only growth performance, but also their energy metabolism. Roque d'Orbcastel et al. (2010) found that increased density also reduced feed intake of *D. labrax*, but Santos et al. (2010) demonstrated that this was partially compensated by a decrease in energy requirements for maintenance. Accordingly, in the present study, differences in growth performance in cobia reared at increased density levels might have been caused by the differential partitioning of energy. It is possible that energy from the diet and/or body reserves was diverted from growth as a response to the elevated metabolic demand caused by stress.

Drip loss is a factor of commercial importance, since elevated levels may affect appearance, juiciness, texture and flavor of the fillet (Kristoffersen et al., 2007). Roth et al. (2006) reported that acute stressed fish had softer texture and an almost three-fold higher drip loss than rested fish. This was probably due to the physical stress of muscle fibrils or connective tissue, which may lead to the release of protease that, in turn, accelerates muscle degeneration. However, drip loss in this study was not affected by density, and values were within those reported for rested fish (Roth et al., 2006). Similar results were recorded for barramundi *Lates calcarifer* (Wilkinson et al., 2008) and king salmon *Oncorhynchus tshawytscha* (Fletcher et al. 2003), in which simulated stress during harvest procedures had no effect on drip loss of their fillets.

Rigor mortis is described as the first post mortem process that has a major influence on the appearance and structure of fish muscle (Berg et al., 1997; Wilkinson et al., 2008). Increased activity and severe stress prior to death may lead to an accelerated breakdown of ATP. Under this condition, fish use anaerobic energy, leading to a rapid drop of muscle pH due to an increased

production of lactic acid. A reduction in muscle pH results in a rapid onset of rigor mortis (Berg et al., 1997; Jerrett et al., 1996; Poli et al., 2005; Bagni et al., 2007). The onset and development of rigor are hence affected by pre-slaughter procedures, and slaughtering methods. They are also influenced by others factors such as species, age and size of the specimen (Berg et al., 1997; Van de Vis et al., 2003; Poli et al., 2005; Bagni et al., 2007; Simitzis et al., 2013). In the present study, stocking density influenced the onset and development of rigor mortis, but the relation between the chronic stressor - stocking density - and flesh quality variables is not clear. No information concerning this topic is currently available on the literature.

Regardless of density level, time from slaughtering to the onset of rigor in cobia may be considered shorter when compared with other fish species exposed to acute stress events. Atlantic salmon (*Salmo solar*), sea bass (*D. labrax*) and sea bream (*S. aurata*) submitted to harvest procedures exhibited full rigor in the range of 6 and 24 h post-mortem (Roth et al., 2006; Bagni et al., 2007; Mørkøre et al., 2008). Nevertheless, our findings are similar to those in barramundi, in which fish submitted to acute stressing agents (air exposure and exercise prior to slaughter) during harvest procedures exhibited a significantly faster onset of rigor, with full rigor being registered 3 h post-mortem. However, as cobia is a tropical fish, differences in rearing temperature for this species and the previously mentioned ones as well as the different methods applied for rigor assessment in the different studies shall also be taken into account. In addition, the slaughter method in this study intended to simulate a commercial slaughter procedure usually applied for cobia. However, this method (Poli et al., 2005; Simitzis et al., 2013) and the one used for rigor measurement (Berg et al., 1997; Skjervold et al. 1999) may have contributed to the earlier onset and development of rigor in juvenile cobia.

In conclusion, stocking densities had no effect on the concentrations of hematological stress indicators, but growth rate and rigor mortis were influenced. Accordingly, juvenile cobia should be maintained at a final biomass up to approximately 15 kg m^{-3} to ensure fish welfare as well a maximal growth rate and an improved final product quality. To our knowledge, this is the first

report assessing the effects of a chronic stressor on both physiological responses and flesh quality parameters in cobia. This may provide information to the advancement of cobia aquaculture, leading to improvements in welfare, final product quality and increased productivity.

5. Acknowledgements

Thanks to Fundação de Amparo à Ciência e Tecnologia de Pernambuco - FACEPE for the provision of a M.Sc. scholarship (IBPG-0694-5.06/10) and a travel grant (AMD-0135-5.04/11) to the first author, and a research grant to R.O. Cavalli (Proc. 0830/5.06-10). Thanks also to Brazil's National Council for Scientific and Technological Development - CNPq for additional financial support (Proc. 559.759/2009-6). R.O. Cavalli is a research fellow of CNPq.

6. References

Ashley, P.J., 2007. Fish welfare: Current issues in aquaculture. Applied Animal Behaviour Science 104, 199-235.

Bagni, M., Civitareale, C., Priori, A., Ballerini, A., Finola, M., Brambilla, G., Marino, G., 2007. Pre-slaughter crowding stress and killing procedures affecting quality and welfare in sea bass (*Dicentrarchus labrax*) and sea bream (*Sparus aurata*). Aquaculture 263, 52–60.

Barton, B.A., 2002. Stress in fishes: a diversity of responses with particular reference to changes in circulating costicosteroids. Integrative and Comparative Biology 42, 517-525.

Barton, B.A., Iwama G.K., 1991. Physiological changes in fish from stress in aquaculture with emphasis on the response and effects of corticosteriods. Annual Review of Fish Diseases 129, 3-26.

Barton, B.A., Ribas, L., Acerete, L., Tort, L., 2005. Effects of chronic confinement on physiological responses of juvenile gilthead sea bream, *Sparus aurata* L., to acute handling. Aquaculture Research 36, 2, 172-179.

Basrur, T.V., Longland, R., Wilkinson, R.J., 2010. Effects of repeated crowding on the stress response and growth performance in Atlantic salmon (*Salmo salar*). Fish Physiology and Biochemistry 36, 445–450.

Benetti, D.D.; O'Hanlon, B.; Riviera, J.A.; Welch, A.W.; Maxey, C.; Orhun, M.R., 2010. Growth rates of cobia (*Rachycentron canadum*) cultured in open ocean submerged cages in the Caribbean. Aquaculture 302, 195-201.

Berg, T., Erikson, U., Nordvedt, T.S., 1997. Rigor mortis assessment of Atlantic salmon (*Salmo salar*) and effects of stress. Journal of Food Science 62, 439–446.

Bianchini, A., Wasielesky, W., Miranda, K.C., 1996. Toxicity of nitrogenous compounds to juveniles of flatfish *Paralichthys orbignyanus*. Bulletin of Environmental Contamination and Toxicology 56, 453-459.

Bito, M., Yamada, K., Mikumo, Y., Amano, K., 1983. Difference in the mode of rigor mortis among some varieties of fish by modified Cutting's method. Bulletin of Tokai Regional Fisheries Research Laboratory 109, 89–93.

Burkey, K., Young, S.P., Smith, T.I.J., Tomasso, J.R., 2007. Low-salinity resistance of juvenile cobias. North American Journal of Aquaculture 69, 271–274.

Canario, A.V.M., Condeca, J., Power, D.M., Ingleton, P. M., 1998. The effect of stocking density on growth in the gilthead sea-bream, *Sparus aurata* (L.). Aquaculture Research 29, 177-181.

Cnaani, A., McLean, E., 2009. Time-course response of cobia (*Rachycentron canadum*) to acute stress. Aquaculture 289, 140–142.

Conte F.S., 2004. Stress and the welfare of cultured fish. Applied Animal Behaviour Science 86, 205-203.

Costas, B., Aragão, C., Dias, J., Afonso, A., Conceição, L.E.C., 2013. Interactive effects of a high-quality protein diet and high stocking density on the stress response and some innate immune parameters of Senegalese sole *Solea senegalensis*. Fish Physiology and Biochemistry. *In press* doi: 10.1007/s10695-013-9770-1

Davis, K.B., 2006. Management of physiological stress in finfish aquaculture. North American Journal of Aquaculture 68, 116–121.

Di Marco, P., Priori, A., Finoia, M.G., Massari, A., Mandich, A., Marino, G., 2008. Physiological responses of European sea bass *Dicentrarchus labrax* to different stocking densities and acute stress challenge. Aquaculture 275, 319–328.

Ellis, T., North, B., Scott, A.P., Bromage, N.R., Porter, M., Gadd, D., 2002. The relationships between stocking density and welfare in farmed rainbow trout. Journal of Fish Biology 61, 3, 493-531.

FAO, 2013. Cultured Aquatic Species Information Programme – *Rachycentron canadum* Available on: www.fao.org/fishery/culturedspecies/Rachycentroncanadum/en [accesse on 29 01 2013].

Fletcher, G.C., Corrigan, V.K., Summers, G., Leonard, M.J., Jerrett, A.R., Black, S.E., 2003. Spoilage of rested harvested king salmon (*Oncorhynchus tshawytscha*). Journal of Food Science 68, 2810–2816.

Herrera, M., Vargas-Chacoff, L., Hachero, I., Ruíz-Jarabo, I., Rodiles A., Navas, J.I. Mancera, J.M., 2009. Physiological responses of juvenile wedge sole *Dicologoglossa cuneata* (Moreau) to high stocking density. Aquaculture Research 40, 790-797.

Iwama G.K., 1998. Stress in fish. Annals of the New York academy of science 851, 304-310.

Jentoft, S., Aastveit A.H., Torjesen, P.A., Andersen, Ø., 2005 Effects of stress on growth, cortisol and glucose levels in non-domesticated Eurasian perch (*Perca fluviatilis*) and domesticated rainbow trout (*Oncorhynchus mykiss*). Comparative Biochemistry and Physiology 141, 353–358.

Kristoffersen, S., Vang, B., Larsen, R., and Olsen, R. L., 2007. Pre-rigor filleting and drip loss from fillets of farmed Atlantic cod (*Gadus morhua* L.). Aquaculture Research 38, 1721-1731.

Liao, I.C.; Huang, T.S.; Tsai, W,-S.; Hsueh, C.M.; Chang, S.L.; Leaño, E.M., 2004. Cobia culture in Taiwan: current status and problems. Aquaculture 237, 155-165.

Liao I.C., Leaño E.D., Hsu C.-Y., Ku C.C., 2007. Marine cage culture of cobia in Taiwan, in: Liao, I.C., Leaño, E.M. (Eds.), Cobia Aquaculture: Research, Development and Commercial Production. Asian Fisheries Society, Manila, Philippines, pp. 131–145.

Mommsen, T.P., Vijayan, M.M., Moon, T.W., 1999. Cortisol in teleosts: dynamics, mechanisms of action, and metabolic regulation. Reviews in Fish Biology and Fisheries 9, 211–268.

Montero D., Izquierdo M., Tort L., Robaina L., Vergara J., 1999. High stocking density produces crowding stress altering some physiological and biochemical parameters in gilthead seabream, *Sparus aurata*, juveniles. Fish Physiology and Biochemistry 20, 53–60.

Mørkøre, T., Mazo, P. I., Tahirovic, V., & Einen, O., 2008. Impact of starvation and handling stress on rigor development and quality of Atlantic salmon (*Salmon salar* L.). Aquaculture 277, 231–238.

Morzel, M., Sohier, D., Van de Vis, H., 2003. Evaluation of slaughtering methods for turbot with respect to animal welfare and flesh quality. Journal of the Science of Food and Agriculture 83, 19–28.

Pickering, A. D., 1993. Growth and stress in fish production. Aquaculture 111, 51-53.

Poli, B.M., Parisi, G., Scappini, F., Zampacavallo, G., 2005. Fish welfare and quality as affected by pre-slaughter and slaughter management. Aquaculture International 13, 29–49.

Procarione, L.S., Barry, T.P., Malison, J.A., 1999. Effects of high rearing densities and loading rates on the growth and stress response of juvenile rainbow trout. North American Journal of Aquaculture 61, 91-96.

Roque d'Orbcastel, E., Lemarié, G., Breuil, G., Petochi, T., Marino, G., Triplet, S., Dutto, G., Fivelstad, S., Coeurdacier, J.L., Blancheton, J.P., 2010. Effects of rearing density on sea bass (*Dicentrarchus labrax*) biological performance, blood parameters and disease resistance in a flow through system. Aquatic Living Resources 23, 109-117.

Roth, B., Slinde, E., & Arildsen, J., 2006. Pre or post mortem muscle activity in Atlantic salmon (Salmo salar). The effect on rigor mortis and the physical properties of flesh. Aquaculture 257, 504–510.

Sammouth, S., d'Orbcastel, E.R., Gasset, E., Lemarié, G., Breuil, G., Marino, G., Coeurdacier, J.L., Fivelstad, S., Blancheton, J.P., 2009. The effect of density on sea bass (*Dicentrarchus labrax*) performance in a tank-based recirculating system. Aquaculture Engineering 40, 2, 72-78.

Sampaio, L.A.; Moreira, C.B.; Miranda-Filho, K.C.; Rombenso, A.N., 2011. Culture of cobia *Rachycentron canadum* (L) in near-shore cages off the Brazilian coast. Aquaculture Research, 42, 832-834.

Santos G.A., Schrama J.W., Mamauag R.E.P., Rombout J.H.W.M., Verreth J.A.J., 2010. Chronic stress impairs performance energy metabolism and welfare indicators in European seabass (*Dicentrarchus labrax*): the combined effects of fish crowding and water quality deterioration. Aquaculture 299, 73–80.

Simitzis, P.E., Tsopelakos, A., Charismiadou, M.A., Batzina, A., Deligeorgis, S.G., Miliou, H., 2013. Comparison of the effects of six stunning/killing procedures on flesh quality of sea bass (*Dicentrarchus labrax*, Linnaeus 1758) and evaluation of clove oil anaesthesia followed by chilling on ice/water slurry for potential implementation in aquaculture. Aquaculture Research. *in press*, doi: 10.1111/are.12120

Skjervold, P.O., Fjæra, S.O., Østby, P.B., 1999. Rigor in Atlantic salmon as affected by crowding stress prior to chilling before slaughter. Aquaculture 175, 93–101.

Sumpter, J. P., 1997. The endocrinology of stress, in Iwama, G. K., Pickering, A.D., Sumpter, J.P., Schreck C.B. (Eds.), Fish stress and health in aquaculture. Inc., New York, pp. 95-118.

Tolussi C.E., Hilsdorf A.W.S., Caneppele D., Moreira R.G., 2010. The effects of stocking density in physiological parameters and growth of the endangered teleost species piabanha, *Brycon insignis* (Steindachner, 1877). Aquaculture 310, 221-228.

Trushenski, J., Schwarz, M., Takeuchi, R., Delbos, B., Sampaio, L.A., 2010. Physiological responses of cobia *Rachycentron canadum* following exposure to low water and air exposure stress challenges. Aquaculture 307, 173-177.

van de Nieuwegiessen, P., Boerlage, A., Verreth, J., Schrama, J., 2008. Assessing the effects of a chronic stressor, stocking density, on welfare indicators of juvenile African catfish, Clarias gariepinus Burchell. Applied Animal Behaviour Science 115, 233–243.

van de Vis, H., Kestin, S., Robb, D., Oehlenschlager, J., Lambooij, B., Munkner, W., Kuhlmann, H., Kloosterboer, K., Tejada, M., Huidobro, A., Ottera, H., Roth, B., Sørensen, N.K., Akse, L., Byrne, H. and Nesvadba, P., 2003. Is humane slaughter of fish possible for industry? Aquaculture Research 34, 211–220.

Wendelaar Bonga S.E., 1997. The stress response in fish. Physiological Reviews 7, 591-625.

Wilkinson, R.J., Paton, N., Porter, M.J.R., 2008. The effects of pre-harvest stress and harvest method on the stress response, rigor onset, muscle pH and drip loss in barramundi (*Lates calcarifer*). Aquaculture 282, 26–32.

Table 1. Mean (± SEM) production performance parameters for cobia (*Rachycentron canadum*) reared at low (LD), medium (MD) and high densities (HD). Means with different letters are significantly different based on one-way ANOVA ($p<0.05$) followed by Duncan's multiple-range test.

	LD	MD	HD	*p*-value
Stocking density (fish L^{-1})	0.07	0.14	0.29	
Initial biomass (kg m^{-3})	3.30 ± 0.02^a	6.67 ± 0.03^b	13.15 ± 0.16^c	<0.001
Final biomass (kg m^{-3})	15.02 ± 0.38^a	25.61 ± 1.5^b	53.94 ± 0.73^c	<0.001
Initial weight (g)	46.27 ± 0.35	46.67 ± 0.24	46.03 ± 0.55	0.56
Final weight (g)	210.27 ± 5.27	192.01 ± 8.57	188.80 ± 2.55	0.09
Survival (%)	100 ± 0	93.3 ± 3.4	100 ± 0	0.36
Weight gain (%)	354.35 ± 7.95^a	311.30 ± 16.32^b	310.13 ± 1.45^b	0.04
SGR (% BW day^{-1})	3.60 ± 0.42^a	3.36 ± 0.09^b	3.36 ± 0.08^b	0.045
FCR	0.98 ± 0.01	1.04 ± 0.05	1.02 ± 0.01	0.42
Feed intake (% BW day^{-1})	3.88 ± 0.04	3.70 ± 0.05	3.70 ± 0.05	0.08
HSI	2.69 ± 0.08	2.57 ± 0.11	2.48 ± 0.05	0.28
K	6.83 ± 0.04	6.70 ± 0.21	6.61 ± 0.07	0.54
Dress-out (%)	36.93 ± 0.45	36.32 ± 1.30	35.92 ± 0.19	0.69

SGR = specific growth rate; BW = body weight; FCR = feed conversion ratio; HSI = hepatosomatic index; K = condition factor.



Figure 1. Mean (± SEM) drip loss (%) from flesh samples obtained from cobia (*Rachycentron canadum*) reared at low (LD), medium (MD) and high stocking densities (HD).



Figure 2. Mean (± SEM) rigor indices (%) for juvenile cobia (*Rachycentron canadum*) raised at low (LD), medium (MD) and high stocking densities (HD). Means with different letters are significantly different within each time point based on one-way ANOVA ($p<0.05$) with post hoc Duncan's multiple-range test.

5. Anexo (Normas para publicação na Aquaculture)

- Introduction

Types of paper

Original Research Papers should report the results of original research. The material should not have been previously published elsewhere. Articles are expected to contribute new information (e.g. novel methods of analysis with added new insights and impacts) to the knowledge base in the field, not just to confirm previously published work.
Review Articles can cover either narrow disciplinary subjects or broad issues requiring interdisciplinary discussion. They should provide objective critical evaluation of a defined subject. Reviews should not consist solely of a summary of published data. Evaluation of the quality of existing data, the status of knowledge, and the research required to advance knowledge of the subject are essential.
Short Communications are used to communicate results which represent a major breakthrough or startling new discovery and which should therefore be published quickly. They should not be used for preliminary results. Papers must contain sufficient data to establish that the research has achieved reliable and significant results.
Technical Papers should present new methods and procedures for either research methodology or culture-related techniques.
The *Letters to the Editor* section is intended to provide a forum for discussion of aquacultural science emanating from material published in the journal.
Contact details for submission

Papers for consideration should be submitted via the electronic submission system mentioned below to the appropriate Section Editor:
Nutrition:
D.M. Gatlin
The Nutrition Section welcomes high quality research papers presenting novel data as well as original reviews on various aspects of aquatic animal nutrition relevant to aquaculture. Manuscripts addressing the following areas of investigation are encouraged:
1) determination of dietary and metabolic requirements for various nutrients by representative aquatic species. Studies may include environmental/stress effects on animal's physiological responses and requirements at different developmental stages;
2) evaluation of novel or established feedstuffs as well as feed processing and manufacturing procedures with digestibility and growth trials. Such studies should provide comprehensive specifications of the process or evaluated ingredients including nutrients, potential anti-nutrients, and contaminants;
3) comparison of nutrient bioavailability from various ingredients or product forms as well as metabolic kinetics of nutrients, food borne anti-nutrients or toxins;
4) identification of key components in natural diets that influence attractability, palatability, metabolism, growth reproduction and/or immunity of cultured organisms;
5) optimization of diet formulations and feeding practices;
6) characterization of the actions of hormones, cytokines and/or components in intracellular signaling pathway(s) that influence nutrient and/or energy utilization.
7) evaluation of diet supplementation strategies to influence animal performance, metabolism, health and/or flesh quality.
Manuscripts concerning other areas of nutrition using novel or advanced methods are also welcome. Please note that in regard to various diet additives such as probiotics, prebiotics, herbal extracts, etc., a very large number of papers have already been published. Therefore, Aquaculture will not continue to accept manuscripts that present initial and preliminary investigations of such additives. Manuscripts addressing these and other feed additives will be accepted for review only if they are of the highest scientific quality and they represent a significant advance in our knowledge of the mechanisms involved in their metabolism. Manuscripts may also be considered if they present clinical efficacy data generated in large-scale trials and economic cost-benefit analysis of these applications.
Aquaculture Production Science:
B.Costa-Pierce
AQUACULTURE PRODUCTION SCIENCE (PS) is one of 5 sections of the international journal AQUACULTURE dedicated to research on improvements and innovations in aquatic food production.
worldwide dissemination of the results of innovative, globally important, scientific research on production methods for aquatic foods from fish, crustaceans, mollusks, amphibians, and all types of aquatic plants. Improvement of production

systems that results in greater efficiencies of resource usage in aquaculture. Effective applications of technologies and methods of aquaculture production for improved stocking regimes, the use of new species and species assemblages, and research on the efficient and sustainable usage of system space with the objective of minimizing resource usage in aquaculture. Investigations to minimize aquaculture wastes and improve water quality, technologies for nutrient recycling in aquaculture ecosystems, and the synergy of aquaculture and other food production systems using methods such as polyculture and integrated aquaculture.

Physiology and Endocrinology:
Fish: A. P. (Tony) Farrell
Invertebrate: J. Benzie
The Physiology Section welcomes high quality papers that present both novel research data and original reviews, on all aspects of the physiology of cultured aquatic animals and plants. Their content must be relevant to solving aquaculture problems.
Submitted manuscripts must have a valid hypothesis or objective, clearly state the relevance to aquaculture, have proper experimental design with appropriate controls and utilize appropriate statistical analysis. Mention of trade names is limited to the main text.
Relevant physiological topics include, but are not limited to:
- Reproductive physiology, including: Endocrine and environmental controls development; Induced ovulation and spermiation; Gamete quality, storage and cryopreservation; control of sex differentiation; Physiology and endocrinology of gynogenetic, triploid and transgenic organisms
- Molecular genetic assessment of physiological processes
- Larval physiology and ontogeny in relation to aquaculture, including metamorphosis, smolting (salmonids) and molting (crustacea)
- Nutritional physiology including endocrine and environmental regulation of growth
- Performance under variable culture conditions, including temperature optima and tolerances; Altered water quality and environmental variables; Stress and disease physiology; Rearing density
- Immunology (physiological studies of probiotics must present statistically valid conclusions)
- Respiratory, muscle and exercise physiology of cultured organisms
- Osmoregulatory physiology and control
- Physiology of harvest and handling techniques, including: Anesthesia and transport; Product and flesh quality; Pigmentation

Genetics:
G. Hulata
The Genetics Section welcomes high-quality research papers presenting novel data, as well as critical reviews, on various aspects of selective breeding, genetics and genomics, so long as the content is relevant to solving aquaculture problems. Please note, however, that Aquaculture will not accept manuscripts dealing with the application of well-described techniques to yet another species, unless the application solves a biological problem important to aquaculture production. Aquaculture will not accept manuscripts dealing with gene cloning, characterizing of microsatellites, species identification using molecular markers, EST papers with small collections, or mapping papers with a small number of markers, unless the papers also deal with solving a biological problem that is relevant to aquaculture production. Where appropriate, linkage maps should include co-dominant markers, such as microsatellite DNA and SNP markers, to enable application to other populations and facilitate comparative mapping. Aquaculture will not accept manuscripts focusing mainly on population genetics studies that are based on RAPD and AFLP markers, since the dominance and multilocus nature of the fingerprints are not suitable for making inferences about population genetic diversity and structure. There may be other journals that are more suitable for manuscripts not meeting these requirements.
Sustainability and Society:
D.C. Little
The Sustainability and Society section of the journal Aquaculture invites articles at the interface of natural and social sciences that address the broader roles of aquaculture in global food security and trade.
Aims and scope of the Sustainability and Society section are the: global dissemination of interdisciplinary knowledge regarding the management of aquatic resources and resulting impacts on people. Interconnections with other sectors of food production; resource management and implications for societal impact. Going beyond a narrow techno-centric focus, towards more holistic analyses of aquaculture within well-defined contexts. Enquiry based on understanding trajectories of change amid the global challenges of climate change and food security. Mixed methods and approaches that incorporate and integrate both social and natural sciences. Relevance for the diverse range of policy makers, practitioners and other stakeholders involved. Articles that take a value chain approach, rather than being wholly production orientated, are encouraged.

Page charges

This journal has no page charges.

Before You Begin

Ethics in publishing

For information on Ethics in publishing and Ethical guidelines for journal publication see http://www.elsevier.com/publishingethics and http://www.elsevier.com/ethicalguidelines.

Policy and ethics

The work described in your article must have been carried out in accordance with *The Code of Ethics of the World Medical Association (Declaration of Helsinki) for experiments involving humans* http://www.wma.net/en/30publications/10policies/b3/index.html; *EU Directive 2010/63/EU for animal experiments* http://ec.europa.eu/environment/chemicals/lab_animals/legislation_en.htm; *Uniform Requirements for manuscripts submitted to Biomedical journals* http://www.icmje.org. This must be stated at an appropriate point in the article.

Conflict of interest

All authors are requested to disclose any actual or potential conflict of interest including any financial, personal or other relationships with other people or organizations within three years of beginning the submitted work that could inappropriately influence, or be perceived to influence, their work. See also http://www.elsevier.com/conflictsofinterest.

Submission declaration and verification

Submission of an article implies that the work described has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see http://www.elsevier.com/postingpolicy), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. To verify originality, your article may be checked by the originality detection service CrossCheck http://www.elsevier.com/editors/plagdetect.

If the manuscript to be submitted was previously rejected by *Aquaculture* or another journal, it is necessary to specify what substantive new work and/or revisions have been included to elevate the manuscript's quality for consideration by *Aquaculture*.

Contributors

Each author is required to declare his or her individual contribution to the article: all authors must have materially participated in the research and/or article preparation, so roles for all authors should be described. The statement that all authors have approved the final article should be true and included in the disclosure.

Changes to authorship

This policy concerns the addition, deletion, or rearrangement of author names in the authorship of accepted manuscripts:

Before the accepted manuscript is published in an online issue: Requests to add or remove an author, or to rearrange the author names, must be sent to the Journal Manager from the corresponding author of the accepted manuscript and must include: (a) the reason the name should be added or removed, or the author names rearranged and (b) written confirmation (e-mail, fax, letter) from all authors that they agree with the addition, removal or rearrangement. In the case of addition or removal of authors, this includes confirmation from the author being added or removed. Requests that are not sent by the corresponding author will be forwarded by the Journal Manager to the corresponding author, who must follow the procedure as described above. Note that: (1) Journal Managers will inform the Journal Editors of any such requests and (2) publication of the accepted manuscript in an online issue is suspended until authorship has been agreed.

After the accepted manuscript is published in an online issue: Any requests to add, delete, or rearrange author names in an article published in an online issue will follow the same policies as noted above and result in a corrigendum.

Copyright

Upon acceptance of an article, authors will be asked to complete a 'Journal Publishing Agreement' (for more information on this and copyright see http://www.elsevier.com/copyright). Acceptance of the agreement will ensure the widest possible dissemination of information. An e-mail will be sent to the corresponding author confirming receipt of the manuscript together with a 'Journal Publishing Agreement' form or a link to the online version of this agreement.

Subscribers may reproduce tables of contents or prepare lists of articles including abstracts for internal circulation within their institutions. Permission of the Publisher is required for resale or distribution outside the institution and for all other derivative works, including compilations and translations (please consult http://www.elsevier.com/permissions). If excerpts from other copyrighted works are included, the author(s) must obtain written permission from the copyright owners and credit the source(s) in the article. Elsevier has preprinted forms for use by authors in these cases: please consult http://www.elsevier.com/permissions.

Retained author rights

As an author you (or your employer or institution) retain certain rights; for details you are referred to: http://www.elsevier.com/authorsrights.

Role of the funding source

You are requested to identify who provided financial support for the conduct of the research and/or preparation of the article and to briefly describe the role of the sponsor(s), if any, in study design; in the collection, analysis and interpretation of data; in the writing of the report; and in the decision to submit the article for publication. If the funding source(s) had no such involvement then this should be stated. Please see http://www.elsevier.com/funding.

Funding body agreements and policies

Elsevier has established agreements and developed policies to allow authors whose articles appear in journals published by Elsevier, to comply with potential manuscript archiving requirements as specified as conditions of their grant awards. To learn more about existing agreements and policies please visit http://www.elsevier.com/fundingbodies.

Open access

This journal does not ordinarily have publication charges; however, authors can now opt to make their articles available to all (including non-subscribers) via the ScienceDirect platform, for which a fee of $3000 applies (for further information on open access see http://www.elsevier.com/about/open-access/open-access-options). Please note that you can only make this choice after receiving notification that your article has been accepted for publication, to avoid any perception of conflict of interest. The fee excludes taxes and other potential costs such as color charges. In some cases, institutions and funding bodies have entered into agreement with Elsevier to meet these fees on behalf of their authors. Details of these agreements are available at http://www.elsevier.com/fundingbodies. Authors of accepted articles, who wish to take advantage of this option, should complete and submit the order form (available at http://www.elsevier.com/locate/openaccessform.pdf). Whatever access option you choose, you retain many rights as an author, including the right to post a revised personal version of your article on your own website. More information can be found here: http://www.elsevier.com/authorsrights.

Language (usage and editing services)

Please write your text in good English (American or British usage is accepted, but not a mixture of these). Authors who feel their English language manuscript may require editing to eliminate possible grammatical or spelling errors and to conform to correct scientific English may wish to use the English Language Editing service available from Elsevier's WebShop http://webshop.elsevier.com/languageediting/ or visit our customer support site http://support.elsevier.com for more information.

Submission

Submission to this journal proceeds totally online and you will be guided stepwise through the creation and uploading of your files. The system automatically converts source files to a single PDF file of the article, which is used in the peer-review process. Please note that even though manuscript source files are converted to PDF files at submission for the review process, these source files are needed for further processing after acceptance. All correspondence, including notification of the Editor's decision and requests for revision, takes place by e-mail removing the need for a paper trail.

Authors should avoid responding by messages received from the system using the 'Reply' button on their e-mail message; this will send the message to the system support and not to the editorial office, and will create unnecessary load of sorting out and forwarding

Please submit your article via http://ees.elsevier.com/aqua/

Referees

Please submit, with the manuscript, the names, addresses and e-mail addresses of three potential referees. Note that the editor retains the sole right to decide whether or not the suggested reviewers are used.

Preparation

Use of wordprocessing software

It is important that the file be saved in the native format of the wordprocessor used. The text should be in single-column format. Keep the layout of the text as simple as possible. Most formatting codes will be removed and replaced on processing the article. In particular, do not use the wordprocessor's options to justify text or to hyphenate words. However, do use bold face, italics, subscripts, superscripts etc. When preparing tables, if you are using a table grid, use only one grid for each individual table and not a grid for each row. If no grid is used, use tabs, not spaces, to align columns. The electronic text should be prepared in a way very similar to that of conventional manuscripts (see also the Guide to Publishing with Elsevier: http://www.elsevier.com/guidepublication). Note that source files of figures, tables and text graphics will be required whether or not you embed your figures in the text. See also the section on Electronic artwork.
To avoid unnecessary errors you are strongly advised to use the 'spell-check' and 'grammar-check' functions of your wordprocessor.

LaTeX

If the LaTeX file is suitable, proofs will be produced without rekeying the text. The article should preferably be written using Elsevier's document class 'elsarticle', or alternatively any of the other recognized classes and formats supported in Elsevier's electronic submissions system, for further information see http://www.elsevier.com/wps/find/authorsview.authors/latex-ees-supported.
The Elsevier 'elsarticle' LaTeX style file package (including detailed instructions for LaTeX preparation) can be obtained from the Quickguide: http://www.elsevier.com/latex. It consists of the file: elsarticle.cls, complete user documentation for the class file, bibliographic style files in various styles, and template files for a quick start.

Article structure

Subdivision - numbered sections

Divide your article into clearly defined and numbered sections. Subsections should be numbered 1.1 (then 1.1.1, 1.1.2, ...), 1.2, etc. (the abstract is not included in section numbering). Use this numbering also for internal cross-referencing: do not just refer to 'the text'. Any subsection may be given a brief heading. Each heading should appear on its own separate line.

Introduction

State the objectives of the work and provide an adequate background, avoiding a detailed literature survey or a summary of the results.

Material and methods

Provide sufficient detail to allow the work to be reproduced. Methods already published should be indicated by a reference: only relevant modifications should be described.

Theory/calculation

A Theory section should extend, not repeat, the background to the article already dealt with in the Introduction and lay the foundation for further work. In contrast, a Calculation section represents a practical development from a theoretical basis.

Results

Results should be clear and concise.

Discussion

This should explore the significance of the results of the work, not repeat them. A combined Results and Discussion section is often appropriate. Avoid extensive citations and discussion of published literature.

Conclusions

The main conclusions of the study may be presented in a short Conclusions section, which may stand alone or form a subsection of a Discussion or Results and Discussion section.

Appendices

If there is more than one appendix, they should be identified as A, B, etc. Formulae and equations in appendices should be given separate numbering: Eq. (A.1), Eq. (A.2), etc.; in a subsequent appendix, Eq. (B.1) and so on. Similarly for tables and figures: Table A.1; Fig. A.1, etc.

Essential title page information

• **Title.** Concise and informative. Titles are often used in information-retrieval systems. Avoid abbreviations and formulae where possible.
• **Author names and affiliations.** Where the family name may be ambiguous (e.g., a double name), please indicate this clearly. Present the authors' affiliation addresses (where the actual work was done) below the names. Indicate all

affiliations with a lower-case superscript letter immediately after the author's name and in front of the appropriate address. Provide the full postal address of each affiliation, including the country name and, if available, the e-mail address of each author.
• **Corresponding author.** Clearly indicate who will handle correspondence at all stages of refereeing and publication, also post-publication. **Ensure that phone numbers (with country and area code) are provided in addition to the e-mail address and the complete postal address. Contact details must be kept up to date by the corresponding author.**
• **Present/permanent address.** If an author has moved since the work described in the article was done, or was visiting at the time, a 'Present address' (or 'Permanent address') may be indicated as a footnote to that author's name. The address at which the author actually did the work must be retained as the main, affiliation address. Superscript Arabic numerals are used for such footnotes.

Abstract

A concise and factual abstract is required. The abstract should state briefly the purpose of the research, the principal results and major conclusions. An abstract is often presented separately from the article, so it must be able to stand alone. For this reason, References should be avoided, but if essential, then cite the author(s) and year(s). Also, non-standard or uncommon abbreviations should be avoided, but if essential they must be defined at their first mention in the abstract itself.

The abstract should be not longer than 400 words.

Keywords

Immediately after the abstract, provide a maximum of 4-6 keywords, using American spelling and avoiding general and plural terms and multiple concepts (avoid, for example, "and", "of"). Be sparing with abbreviations: only abbreviations firmly established in the field may be eligible. These keywords will be used for indexing purposes.

Abbreviations

Define abbreviations that are not standard in this field in a footnote to be placed on the first page of the article. Such abbreviations that are unavoidable in the abstract must be defined at their first mention there, as well as in the footnote. Ensure consistency of abbreviations throughout the article.

Acknowledgements

Collate acknowledgements in a separate section at the end of the article before the references and do not, therefore, include them on the title page, as a footnote to the title or otherwise. List here those individuals who provided help during the research (e.g., providing language help, writing assistance or proof reading the article, etc.).

Nomenclature and units

Follow internationally accepted rules and conventions: use the international system of units (SI). If other quantities are mentioned, give their equivalent in SI. You are urged to consult IUPAC: Nomenclature of Organic Chemistry: http://www.iupac.org/ for further information.

1. Authors and editors are, by general agreement, obliged to accept the rules governing biological nomenclature, as laid down in the International Code of Botanical Nomenclature, the International Code of Nomenclature of Bacteria, and the International Code of Zoological Nomenclature.
2. All biota (crops, plants, insects, birds, mammals, etc.) should be identified by their scientific names when the English term is first used, with the exception of common domestic animals.
3. All biocides and other organic compounds must be identified by their Geneva names when first used in the text. Active ingredients of all formulations should be likewise identified.
4. For chemical nomenclature, the conventions of the International Union of Pure and Applied Chemistry and the official recommendations of the IUPAC IUB Combined Commission on Biochemical Nomenclature should be followed.

Database linking

Elsevier encourages authors to connect articles with external databases, giving their readers one-click access to relevant databases that help to build a better understanding of the described research. Please refer to relevant database identifiers using the following format in your article: Database: xxxx (e.g., TAIR: AT1G01020; CCDC: 734053; PDB: 1XFN). See http://www.elsevier.com/databaselinking for more information and a full list of supported databases.

DNA sequences and GenBank Accession numbers. Many Elsevier journals cite "gene accession numbers" in their running text and footnotes. Gene accession numbers refer to genes or DNA sequences about which further information can be found in the databases at the National Center for Biotechnical Information (NCBI) at the National Library of

Medicine. Authors are encouraged to check accession numbers used very carefully. **An error in a letter or number can result in a dead link.** Note that in the final version of the electronic copy, the accession number text will be linked to the appropriate source in the NCBI databases enabling readers to go directly to that source from the article.
Example 1: "GenBank accession nos. **AI631510, AI631511, AI632198,** and **BF223228**, a B-cell tumor from a chronic lymphatic leukemia (GenBank accession no. BE675048), and a T-cell lymphoma (GenBank accession no. **AA361117**)".
Authors are encouraged to check accession numbers used very carefully. An error in a letter or number can result in a dead link.
In the final version of the printed article, the accession number text will not appear bold or underlined (see Example 2 below).
Example 2: "GenBank accession nos. AI631510, AI631511, AI632198, and BF223228), a B-cell tumor from a chronic lymphatic leukemia (GenBank accession no. BE675048), and a T-cell lymphoma (GenBank accession no. AA361117)".
In the final version of the electronic copy, the accession number text will be linked to the appropriate source in the NCBI databases enabling readers to go directly to that source from the article (see Example 3 below).
Example 3: "GenBank accession nos. AI631510, AI631511, AI632198, and BF223228), a B-cell tumor from a chronic lymphatic leukemia (GenBank accession no. BE675048), and a T-cell lymphoma (GenBank accession no. AA361117)".
Math formulae

Present simple formulae in the line of normal text where possible and use the solidus (/) instead of a horizontal line for small fractional terms, e.g., X/Y. In principle, variables are to be presented in italics. Powers of e are often more conveniently denoted by exp. Number consecutively any equations that have to be displayed separately from the text (if referred to explicitly in the text).

Give the meaning of all symbols immediately after the equation in which they are first used. In chemical formulae, valence of ions should be given as, e.g. Ca^{2+} and not Ca^{++}. Isotope numbers should precede the symbols, e.g., ^{18}O. The repeated writing of chemical formulae in the text is to be avoided where reasonably possible; instead, the name of the compound should be given in full. Exceptions may be made in the case of a very long name occurring very frequently or in the case of a compound being described as the end product of a gravimetric determination (e.g., phosphate as P_2O_5).
Footnotes

Footnotes should be used sparingly. Number them consecutively throughout the article, using superscript Arabic numbers. Many wordprocessors build footnotes into the text, and this feature may be used. Should this not be the case, indicate the position of footnotes in the text and present the footnotes themselves separately at the end of the article. Do not include footnotes in the Reference list.
Table footnotes
Indicate each footnote in a table with a superscript lowercase letter.
Artwork
Electronic artwork
General points
• Make sure you use uniform lettering and sizing of your original artwork.
• Embed the used fonts if the application provides that option.
• Aim to use the following fonts in your illustrations: Arial, Courier, Times New Roman, Symbol, or use fonts that look similar.
• Number the illustrations according to their sequence in the text.
• Use a logical naming convention for your artwork files.
• Provide captions to illustrations separately.
• Size the illustrations close to the desired dimensions of the printed version.
• Submit each illustration as a separate file.
A detailed guide on electronic artwork is available on our website:
http://www.elsevier.com/artworkinstructions
You are urged to visit this site; some excerpts from the detailed information are given here.
Formats
If your electronic artwork is created in a Microsoft Office application (Word, PowerPoint, Excel) then please supply 'as is' in the native document format.
Regardless of the application used other than Microsoft Office, when your electronic artwork is finalized, please 'Save as' or convert the images to one of the following formats (note the resolution requirements for line drawings, halftones, and line/halftone combinations given below):
EPS (or PDF): Vector drawings, embed all used fonts.
TIFF (or JPEG): Color or grayscale photographs (halftones), keep to a minimum of 300 dpi.
TIFF (or JPEG): Bitmapped (pure black & white pixels) line drawings, keep to a minimum of 1000 dpi.
TIFF (or JPEG): Combinations bitmapped line/half-tone (color or grayscale), keep to a minimum of 500 dpi.

Please do not:
• Supply files that are optimized for screen use (e.g., GIF, BMP, PICT, WPG); these typically have a low number of pixels and limited set of colors;
• Supply files that are too low in resolution;
• Submit graphics that are disproportionately large for the content.

Color artwork

Please make sure that artwork files are in an acceptable format (TIFF, EPS or MS Office files) and with the correct resolution. If, together with your accepted article, you submit usable color figures then Elsevier will ensure, at no additional charge, that these figures will appear in color on the Web (e.g., ScienceDirect and other sites) regardless of whether or not these illustrations are reproduced in color in the printed version. **For color reproduction in print, you will receive information regarding the costs from Elsevier after receipt of your accepted article.** Please indicate your preference for color: in print or on the Web only. For further information on the preparation of electronic artwork, please see http://www.elsevier.com/artworkinstructions.

Please note: Because of technical complications which can arise by converting color figures to 'gray scale' (for the printed version should you not opt for color in print) please submit in addition usable black and white versions of all the color illustrations.

Figure captions

Ensure that each illustration has a caption. Supply captions separately, not attached to the figure. A caption should comprise a brief title (**not** on the figure itself) and a description of the illustration. Keep text in the illustrations themselves to a minimum but explain all symbols and abbreviations used.

Text graphics

Text graphics may be embedded in the text at the appropriate position. See further under Electronic artwork.

Tables

Number tables consecutively in accordance with their appearance in the text. Place footnotes to tables below the table body and indicate them with superscript lowercase letters. Avoid vertical rules. Be sparing in the use of tables and ensure that the data presented in tables do not duplicate results described elsewhere in the article.

References

Citation in text

Please ensure that every reference cited in the text is also present in the reference list (and vice versa). Any references cited in the abstract must be given in full. Unpublished results and personal communications are not recommended in the reference list, but may be mentioned in the text. If these references are included in the reference list they should follow the standard reference style of the journal and should include a substitution of the publication date with either 'Unpublished results' or 'Personal communication'. Citation of a reference as 'in press' implies that the item has been accepted for publication.

Web references

As a minimum, the full URL should be given and the date when the reference was last accessed. Any further information, if known (DOI, author names, dates, reference to a source publication, etc.), should also be given. Web references can be listed separately (e.g., after the reference list) under a different heading if desired, or can be included in the reference list.

References in a special issue

Please ensure that the words 'this issue' are added to any references in the list (and any citations in the text) to other articles in the same Special Issue.

Reference management software

This journal has standard templates available in key reference management packages EndNote (http://www.endnote.com/support/enstyles.asp) and Reference Manager (http://refman.com/support/rmstyles.asp). Using plug-ins to wordprocessing packages, authors only need to select the appropriate journal template when preparing their article and the list of references and citations to these will be formatted according to the journal style which is described below.

Reference style

Text: All citations in the text should refer to:
1. *Single author:* the author's name (without initials, unless there is ambiguity) and the year of publication;
2. *Two authors:* both authors' names and the year of publication;
3. *Three or more authors:* first author's name followed by 'et al.' and the year of publication.

Citations may be made directly (or parenthetically). Groups of references should be listed first alphabetically, then chronologically.

Examples: 'as demonstrated (Allan, 2000a, 2000b, 1999; Allan and Jones, 1999). Kramer et al. (2010) have recently shown ....'

List: References should be arranged first alphabetically and then further sorted chronologically if necessary. More than one reference from the same author(s) in the same year must be identified by the letters 'a', 'b', 'c', etc., placed after the

year of publication.
Examples:
Reference to a journal publication:
Van der Geer, J., Hanraads, J.A.J., Lupton, R.A., 2010. The art of writing a scientific article. J. Sci. Commun. 163, 51–59.
Reference to a book:
Strunk Jr., W., White, E.B., 2000. The Elements of Style, fourth ed. Longman, New York.
Reference to a chapter in an edited book:
Mettam, G.R., Adams, L.B., 2009. How to prepare an electronic version of your article, in: Jones, B.S., Smith , R.Z. (Eds.), Introduction to the Electronic Age. E-Publishing Inc., New York, pp. 281–304.

Journal Abbreviations Source

Define abbreviations that are not standard in this field at their first occurrence in the article: in the abstract but also in the main text after it. Ensure consistency of abbreviations throughout the article.

Video data

Elsevier accepts video material and animation sequences to support and enhance your scientific research. Authors who have video or animation files that they wish to submit with their article are strongly encouraged to include links to these within the body of the article. This can be done in the same way as a figure or table by referring to the video or animation content and noting in the body text where it should be placed. All submitted files should be properly labeled so that they directly relate to the video file's content. In order to ensure that your video or animation material is directly usable, please provide the files in one of our recommended file formats with a preferred maximum size of 50 MB. Video and animation files supplied will be published online in the electronic version of your article in Elsevier Web products, including ScienceDirect: http://www.sciencedirect.com. Please supply 'stills' with your files: you can choose any frame from the video or animation or make a separate image. These will be used instead of standard icons and will personalize the link to your video data. For more detailed instructions please visit our video instruction pages at http://www.elsevier.com/artworkinstructions. Note: since video and animation cannot be embedded in the print version of the journal, please provide text for both the electronic and the print version for the portions of the article that refer to this content.

Supplementary data

Elsevier accepts electronic supplementary material to support and enhance your scientific research. Supplementary files offer the author additional possibilities to publish supporting applications, high-resolution images, background datasets, sound clips and more. Supplementary files supplied will be published online alongside the electronic version of your article in Elsevier Web products, including ScienceDirect: http://www.sciencedirect.com. In order to ensure that your submitted material is directly usable, please provide the data in one of our recommended file formats. Authors should submit the material in electronic format together with the article and supply a concise and descriptive caption for each file. For more detailed instructions please visit our artwork instruction pages at http://www.elsevier.com/artworkinstructions.

Submission checklist

The following list will be useful during the final checking of an article prior to sending it to the journal for review. Please consult this Guide for Authors for further details of any item.

Ensure that the following items are present:

One author has been designated as the corresponding author with contact details:
• E-mail address
• Full postal address
• Phone numbers
All necessary files have been uploaded, and contain:
• Keywords
• All figure captions
• All tables (including title, description, footnotes)
Further considerations
• Manuscript has been 'spell-checked' and 'grammar-checked'
• References are in the correct format for this journal
• All references mentioned in the Reference list are cited in the text, and vice versa
• Permission has been obtained for use of copyrighted material from other sources (including the Web)
• Color figures are clearly marked as being intended for color reproduction on the Web (free of charge) and in print, or to be reproduced in color on the Web (free of charge) and in black-and-white in print
• If only color on the Web is required, black-and-white versions of the figures are also supplied for printing purposes
For any further information please visit our customer support site at http://support.elsevier.com.

After Acceptance

Use of the Digital Object Identifier

The Digital Object Identifier (DOI) may be used to cite and link to electronic documents. The DOI consists of a unique alpha-numeric character string which is assigned to a document by the publisher upon the initial electronic publication. The assigned DOI never changes. Therefore, it is an ideal medium for citing a document, particularly 'Articles in press' because they have not yet received their full bibliographic information. Example of a correctly given DOI (in URL format; here an article in the journal *Physics Letters B*):
http://dx.doi.org/10.1016/j.physletb.2010.09.059
When you use a DOI to create links to documents on the web, the DOIs are guaranteed never to change.

Proofs

One set of page proofs (as PDF files) will be sent by e-mail to the corresponding author (if we do not have an e-mail address then paper proofs will be sent by post) or, a link will be provided in the e-mail so that authors can download the files themselves. Elsevier now provides authors with PDF proofs which can be annotated; for this you will need to download Adobe Reader version 7 (or higher) available free from http://get.adobe.com/reader. Instructions on how to annotate PDF files will accompany the proofs (also given online). The exact system requirements are given at the Adobe site: http://www.adobe.com/products/reader/tech-specs.html.
If you do not wish to use the PDF annotations function, you may list the corrections (including replies to the Query Form) and return them to Elsevier in an e-mail. Please list your corrections quoting line number. If, for any reason, this is not possible, then mark the corrections and any other comments (including replies to the Query Form) on a printout of your proof and return by fax, or scan the pages and e-mail, or by post. Please use this proof only for checking the typesetting, editing, completeness and correctness of the text, tables and figures. Significant changes to the article as accepted for publication will only be considered at this stage with permission from the Editor. We will do everything possible to get your article published quickly and accurately – please let us have all your corrections within 48 hours. It is important to ensure that all corrections are sent back to us in one communication: please check carefully before replying, as inclusion of any subsequent corrections cannot be guaranteed. Proofreading is solely your responsibility. Note that Elsevier may proceed with the publication of your article if no response is received.

Offprints

The corresponding author, at no cost, will be provided with a PDF file of the article via e-mail (the PDF file is a watermarked version of the published article and includes a cover sheet with the journal cover image and a disclaimer outlining the terms and conditions of use). For an extra charge, paper offprints can be ordered via the offprint order form which is sent once the article is accepted for publication. Both corresponding and co-authors may order offprints at any time via Elsevier's WebShop (http://webshop.elsevier.com/myarticleservices/offprints). Authors requiring printed copies of multiple articles may use Elsevier WebShop's 'Create Your Own Book' service to collate multiple articles within a single cover (http://webshop.elsevier.com/myarticleservices/offprints/myarticlesservices/booklets).

Author Inquiries

For inquiries relating to the submission of articles (including electronic submission) please visit this journal's homepage. For detailed instructions on the preparation of electronic artwork, please visit http://www.elsevier.com/artworkinstructions. Contact details for questions arising after acceptance of an article, especially those relating to proofs, will be provided by the publisher. You can track accepted articles at http://www.elsevier.com/trackarticle. You can also check our Author FAQs at http://www.elsevier.com/authorFAQ and/or contact Customer Support via http://support.elsevier.com.